

People's Democratic Republic of Algeria  
وزارة التعليم العالي والبحث العلمي  
Ministry of Higher Education and Scientific Research  
جامعة ابن خلدون - تيارت  
University Ibn Khaldoun – Tiaret-



Faculty of Material Sciences

كلية علوم المادة

Chemistry Department

قسم الكيمياء

## Memoir

*Presented by:*

Ms. ABDERRHIME IMENE

Ms. BENAOUA CHAHINEZ HIBA

*To obtain the diploma of*

*Master's degree*

*Startup project 1275*

**Stream: Chemistry**

**Specialization: Organic Chemistry**

**Subject:**

**Evaluation of the biological activity of galenic forms  
prepared from medicinal plants**

**Defended on: 02/07/2024**

**Before the jury:**

**M<sup>r</sup>. DAHO Bouabdellah**

**M<sup>me</sup>. MABREK Houria**

**M<sup>me</sup>. ABDELMALEK Ilham**

**M<sup>lle</sup>. BENNABI Lamia**

**M<sup>me</sup>. TALBI Khadidja**

**M<sup>r</sup>. MOUMEN Taqiyeddine**

**President**

**Examiner**

**Promoter**

**Co-promoter**

**Pharmacist**

**University**

**incubator**

**representative**

**CMB at Tiaret - UNIV**

**AMA at Tiaret - UNIV**

**CMA at Tiaret - UNIV**

**CMA at Ain Timouchent - UNIV**

**CMA at Tiaret - UNIV**

**Academic year: 2023/2024**

# Thanks

*First and foremost, we wish to express our gratitude to God, the Almighty and Merciful, for giving us the c*

*ourage and patience to complete this modest work, and for protecting and keeping us in good health during this delicate period.*

*Our thanks then go to our supervisor, Madame. I. Abdelmalek, who has continuously advised, encouraged, and supported us. We are deeply grateful for the experience she has shared with us and for the valuable time she has dedicated to guiding this work.*

*We also extend our warm and our heartfelt thanks to our co-supervisor, Miss L. Bennabi for her efforts in this work despite the long distances and her presence with us for guidance and advice, so thank you very much.*

*We extend our sincere thanks to the members of the jury, M<sup>me</sup> H. Mebrek, AMA. Mr B. Dahou CMB and Mr. Moumen CMA from Ibn Khaldoun University, for agreeing to evaluate this work.*

*We wish to express our sincere gratitude to the entire staff of the various laboratories:*

- From the Chemistry Laboratory: M<sup>r</sup>F. Babou, M<sup>r</sup>A. Larbi, and Mrs. S. Kouadria.*
- From the Synthesis and Catalysis Laboratory: Ms. N. Sahnoune. M<sup>r</sup>. E. Belarbi.*
- And finally, from the Farm Animal Reproduction Laboratory: Miss R. Bouteldja and M<sup>r</sup> R. Doucene, for providing us with all the necessary equipment and products. Thank you for your availability, generosity, and assistance.*

*We express our gratitude to all the teachers who have contributed to our education.*

*We do not forget to thank our colleagues N. Benabdelah for the support and help specially in our work and our groupe” futures chimie “, for their support, encouragement, unforgettable moments of fun, joy, and sadness we shared during all these years of study.*

*Words in any language will never be sufficient to describe our gratitude to our dear parents, brothers, and sisters. Your love and presence have always illuminated and guided our steps.*

*This manuscript expresses our gratitude and affection to all those who contributed, directly or indirectly, to its completion and who kindly made this work a very rewarding experience.*

## اهداء

"بسم هلا خالقي وميسر اموري وعصمة امري لك كل الحمد والامتنان منال"

من قال ان لها .. نالها

لم تكن الرحلة قصيرة وال ذنبغي لها ان نكون لم يكن الحلم قريبا وال الطريق محفوفا بالتسهيلات لكنني فعلتها ونالها  
الى من كلفه هلا بالهبة والوقار.. الى من احمل اسمه بكل فخر.. الى من حصد الشواك عن دربي ليمهد لي طريق العلم  
بعد بعد فضل هلا ما ان فيه يعود الى .. ابي- الرجل الذي سعى طوال حياته لكي يكون أفضل منه

- ابي الغالي -

الى م الكي في الحياة الى معنى الحب والى معنى الحنان والنفاري الى بسممة الحياة وسر الوجود الى من كان دعائه سر  
الى الم ولخت والصديقة داعمي الأول ووجته الذي استمد منها القوة امي الحبيبة صديقة داعمي الأول ووجتي التي استمد منها القوة.

الى مصدر قوتي الداعمين السانين رضى الصلابة وجراري المنين الى من وهب علي تعالي.. ويدكروني بمدى قوتي  
واستطاعتي الذين ال يحبونني ويؤمنوا بشجاعتني مهما ضربت ورنخت واقفين خلفي مثل ظال مهما كثرة تخبطاتي من  
بذلوا جهدا في مساعدي واداروا عونا وسندا الى الشروع التي تهر طريق  
واستطاعتي للذين لا يحبطوني ويؤمنوا بشجاعتني مهما ضربت ورنخت واقفين خلفي.. مثل ظالا مهما كثرة تخبطاتي من

بذلوا جهدا واخواني امينة وخطرة الزهرع منال تنير طريقني

-اخواني واخواني - امينة - مخطرة - زهرة - منال -

ح والذين شاركوني خطوات هذا الطريق الى من هوروا نعب الطريق الى من شجعوني على امثابرة التمام  
ولا انمي رفقاء الروح والذين شاركوني خطوات هذا الطريق الى من شجعوني على المثابرة  
مسيرتي الى رفقاء السنين ممتنة لكم لالكم باسمه

لإتمام مسوتي الى رفقاء السنين ممتنة لكم كلكم باسمه  
شيماء صبرين خولة امينة كريمة عبدالعزير امين احمد

-شيماء-صبرين -خولة -امينة -كريمة -عبدالعزير -امين -احمد-

الى سندي ومسندي صديقتي واختي الى من شارككتني خطوات هذا الطريق بكل عقباته ونجاحاته خيرة ايامي وصفوتها  
الى سندي ومسندي صديقتي واختي ... الى من شارككتني خطوات هذا الطريق بكل عقباته ونجاحاته.. خيرة ايامي وصفوتها

زميلتي في هذا العمل غاليتي

ايمان

-ايمان-

لله الشكر كله ان وفقني لهذه اللحظة فالحمد لله رب العالمين والصلاة والسلام على نبيه الكريم



CHAHINEZ HIBA



## اهداء

"بسم هلا خالقي وميسر اموري وعصرة اجري لك كل الحمد والمنة ان"

من قال ان الهاء "ناله"

لم تكن الرحلة قصيرة والينبغي لها ان تكون لم تكن الحلم قريبا وال الطريق حفوفا بالنسبة يلات لكنني فعلتها وبلتها.

الى رجل الكفاح من دعمني بالحدود واعطاني بال مقابل... الى من زين اسمي بأجمل اللقاب وحصد الشواك عن دربي ليهدد لي طريق العلم بعد فضل هلا.. ما انا فيه يعود الى

- ابي الغالي -

الى ملكي الطاهر وقوتي بعد هلا داعمني البداية التي احتضرتني بقله اقبل بديه وسهلت لي الشدائد بدعائه اهديك هذا الإنجاز الذي لو ان نضحتك هلا لولا تضحيته كان له وجود ممتنة ان هلا اصطفك لي من البشر اما يا خير سند وعوض سند وعوض

- امي الحنوننة -

الى من قبل فيهم (سَنَسَ دَعَضَ دَكَ يَا خِي كَ)

الى أجمل الهدايا التي قومتها لي عائلتي... من مد يده دون ليل والليل وقت ضعفي - اخي -

الى اميرات ابي ورفيقات دربي... الى من آمن بقدراتي وأمان أيامي.. الى من تذكرني بقوتي وتقفن خلفي كظلي

- أخواتي -

الى رفقاء الخطوة الأولى والخطوة ما قبل لخرة... الى من كانوا لي السروات العجاف سحابا ممطرا.. رفقاء الروح والحرب والسنين

- شرياء - صبرين - خولة - امينة - كريمة - عبد العزيز - امين - احمد -

الى من نانس نفسي لفسها... الى من توافقني الفكر والطبع والقيم.. ولن الصداقة حظور رزق اشكر هلا ان رزقني اياك.. صديقتي

وزميلتي في هذا العمل

- شهيناز -

لله الشكر كله ان وفقني لهذه اللحظة فالحمد لله رب العالمين والصلاة والسلام على نبيه الكريم



IMENE

## المخلص

يركز عملنا على استخلاص المركبات العلاجية من النباتات، وصياغة العلاجات الفعّالة على المستخلصات النباتية وتقييم إمكاناتها العلاجية. تمّ تطوير طرق الاستخلاص وتحضير المستخلص، بالإضافة إلى التجارب المخبرية والحيوية لتقييم النشاط المضاد للأكسدة والمضاد للالتهابات والقدرة على تعزيز نمو الشعر في الفئران. أظهرت النتائج أن مستخلص الميثان يُظهر نشاطًا كبيرًا مضادًا للأكسدة، بينما يُظهر مستخلص البابونج نشاطًا أول وضوحًا مضادًا للالتهابات. تم اختبار التركيبات الرجولية التي تم تحضيرها على الفئران، وأظهرت تأثيرات إيجابية على اللثة ونمو الشعر.

## Résumé

القامات النباتية: تحارب على الجسم الحي، نشاط مضاد الجراثيم، مستخلصات الميثان والبابونج  
Notre travail concentre sur l'extraction de composés thérapeutiques à partir de plantes,

la formulation de remèdes à base d'extraits de plantes et l'évaluation de leur potentiel thérapeutique. Des méthodes d'extraction et de préparation d'extraits sont détaillées, ainsi que des tests in vitro et in vivo pour évaluer l'activité antioxydante, anti-inflammatoire et la capacité de favoriser la croissance des poils chez les rats. Les résultats montrent que l'extrait de Thymelaea présente une activité antioxydante significative, tandis que l'extrait de Chamomile montre une activité anti-inflammatoire moins prononcée. Les formulations galéniques préparées ont été testées sur des rats, démontrant des effets positifs sur l'inflammation et la croissance des poils.

**Mots-clés :** Méthane et extraits de camomille, expériences in vivo, activité antibactérienne.

## Abstract

Our work focuses on the extraction of therapeutic compounds from plants, the formulation of plant extract-based remedies and the evaluation of their therapeutic potential. Extraction and extract preparation methods are detailed, as well as in vitro and in vivo tests to assess antioxidant and anti-inflammatory activity and the ability to promote hair growth in rats. The results show that Thymelaea extract exhibits significant antioxidant activity, while Chamomile extract shows less pronounced anti-inflammatory activity. The galenic formulations prepared were tested on rats, demonstrating positive effects on inflammation and hair growth.

**Keywords:** Methane and chamomile extracts, in vivo experiments, antibacterial activity

## ***Summary***

|   |    |
|---|----|
| <i>General Introduction</i> .....                   | 1  |
| I.1. INTRODUCTION .....                             | 2  |
| I.2. DEFINITION OF MEDICINAL PLANTS .....           | 2  |
| I.3. THE BENEFITS OF MEDICINAL PLANT.....           | 2  |
| I.4. PRINCIPAL INGREDIENTS OF MEDICINAL PLANTS..... | 4  |
| I.5. THE STUDIED PLANTS .....                       | 4  |
| <i>I.5.1. Thymelaea hirsuta</i> .....               | 5  |
| <i>I.5.2. Chamomile</i> .....                       | 9  |
| <i>I.5.3. Curcuma longa</i> .....                   | 13 |
| II.1. INTRODUCTION.....                             | 18 |
| II.2. MATERIALS AND METHODS .....                   | 18 |
| II.2.1. Chemical products .....                     | 18 |
| II.2.2. Vegetal materials .....                     | 18 |
| II.2.3. Methodology of work .....                   | 19 |
| II.2.4. Preparation of the extract.....             | 23 |
| II.3. QUALITATIVE ANALYSIS .....                    | 23 |
| II.4. PHYTOCHEMICAL SCREENING .....                 | 24 |
| II.5. QUANTITATIVE ANALYSIS OF EXTRACTS.....        | 25 |
| II.5.1. Dosage of total polyphenols .....           | 25 |
| II.5.2. Determination of total flavonoids.....      | 26 |
| II.5.3. Dosage of condensed tannins .....           | 27 |
| II.6. GALENIC FORMS FORMULATION.....                | 28 |
| II.7. QUALITY CONTROL FOR GALENIC FORMS.....        | 30 |
| II.8. DRUG RELEASE OF GALENIC FORMS .....           | 30 |
| II.8.1. Operating procedure .....                   | 30 |
| II.8.2. Principe active analysis .....              | 32 |
| II.9. BIOLOGICAL STUDY .....                        | 35 |

|  |    |
|--|----|
| II.10. MINIMAL CONCENTRATION OF INHIBITORY .....                   | 38 |
| II.10.1. Principe of the method.....                               | 38 |
| II.10.2. Protocol of the MCI Determination.....                    | 39 |
| II.11. ANTIOXIDANT ACTIVITY .....                                  | 40 |
| II.11.1. Principe of the method .....                              | 40 |
| II.11.2. Antioxidant experiment .....                              | 41 |
| II.11.3. IC50 estimation .....                                     | 42 |
| II.12. ANTI-INFLAMMATORY ACTIVITY.....                             | 42 |
| II.13. IN VIVO TESTS .....   | 43 |
| III.1. INTRODUCTION .....  | 47 |
| III.2. CHARACTERISATION OF EXTRACTS.....                           | 47 |
| III.2.1 Physico-chemical characterization of extracts product..... | 47 |
| III.2.2. Phytochemical analysis .....                              | 52 |
| III.2.3. Quantitative analysis .....                               | 53 |
| III.3. CHARACTERISATION OF GALINICS FORMS .....                    | 55 |
| III.4. RELEASE STUDY .....   | 57 |
| III.5. BACTERIAL TEST.....   | 61 |
| III.6. MINIMAL CONCENTRATION INHIBITORY .....                      | 64 |
| III.7. ANTIOXYDANT ACTIVITY.....                                   | 67 |
| III.8. ANTIINFLAMMATORY ACTIVITY .....                             | 68 |
| III.9. IN VIVO TEST .....  | 68 |
| <i>General conclusion</i> .....                                    | 72 |

## LIST OF ABBREVIATIONS

|                |  |
|----------------|--|
| Abs            | Absorption   |
| ATCC           | American Type Culture Collection.                                      |
| C              | Concentration of the solution in mole/L                                |
| °C             | Celcius degree   |
| Cam            | Matricaria Chamomile   |
| Candida        | <i>Candida albicans</i>  |
| Cur            | Curcuma  |
| DPPH           | 2,2-diphenyl-1-picrylhydrazyl  |
| DMSO           | Di Methyl Sulfoxide  |
| E.coli         | <i>Escherichia coli</i>  |
| I max          | Maximal Intensity  |
| I <sub>0</sub> | Initial Intensity  |
| l              | Length of quartz cell (1 cm )  |
| L1             | Cream of Curcuma 1 %   |
| L1.0           | Blank of L1  |
| L2             | Cream of curcuma 1% + zinc oxyde                                       |
| L2.0           | Blank of L2  |
| L3             | Cream of Chamomile 1 %   |
| L3.0           | Blank of L3  |
| L4             | Cream of Chamomile 1.5%  |
| L4.0           | Blank of L4  |
| L5             | Cream of curcumin 2%+1% Chamomile                                      |
| L5.0           | Blank of L5  |
| L6             | Cream of curcumin 1%+2% Chamomile                                      |
| L6.0           | Blanc of L6  |
| L7             | Cream of thymelaea 1%  |
| L7.0           | Blank of L7  |
| m              | Mass of dry extract  |
| MIC            | Minimal Inhibitory Concentration                                       |
| nm             | Nano miter   |
| pH             | Hydrogen potential   |
| Pseudo         | <i>Pseudomonas aeruginosa</i>  |
| rfm            | Rote for minute  |
| Staph          | <i>Staphylococcus aureus</i>   |
| TC             | Tannin Content   |
| TFC            | Total Flavonoid Content  |
| TPC            | Total Polyphenols Content  |
| UV             | Ultra- Violet  |
| V              | Extract volume   |
| VF             | Volume of residues solution in the flacon following each precipitation |
| Vf             | Volume of solution after dilution                                      |
| Vp             | Taken Volume   |
| ε              | the specific absorption coefficient                                    |
| μL             | Micro liter  |



# Tables List

## *Bibliography part*

|  |    |
|--|----|
| <b>Table I.1:</b> the classification of <i>Thymelaea hirsuta</i> ..... | 7  |
| <b>Table I.2:</b> The scientific classification of Chamomile .....     | 11 |

## *Experimental part*

|   |    |
|---|----|
| <b>Table II.1:</b> chemical products .....                                      | 18 |
| <b>Table II.2:</b> The composition of L1 .....                                  | 28 |
| <b>Table II.3:</b> The composition of L2.....                                   | 29 |
| <b>Table II.4:</b> The composition of L3.....                                   | 29 |
| <b>Table II.5:</b> The composition of L4.....                                   | 29 |
| <b>Table II.6:</b> The composition of L5.....                                   | 29 |
| <b>Table II.7:</b> The composition of L6.....                                   | 29 |
| <b>Table II.8:</b> The composition of L7.....                                   | 29 |
| <b>Table II.9:</b> The composition of Shampoo .....                             | 29 |
| <b>Table II.10:</b> The tested bacterial stains .....                           | 36 |
| <b>Table II.11:</b> Description of rat's group of the first experience .....    | 45 |
| <b>Table II.12:</b> Description of rat's groups for the second experience ..... | 46 |

## *Results and discussion part*

|  |    |
|--|----|
| <b>Table III.1:</b> The calculated yield .....   | 47 |
| <b>Table III.2:</b> The solubility's test .....  | 48 |
| <b>Table III.3:</b> Characteristic bands of extract plants .....                               | 51 |
| <b>Table III.4:</b> Result of phytochemical tests.....   | 52 |
| <b>Table III.5:</b> Result of centrifugation of creams .....                                   | 55 |
| <b>Table III.6:</b> The galenic forms pH.....  | 56 |
| <b>Table III.7:</b> Result of antibacterial activity of Thy extract with the method of MIC ... | 65 |
| <b>Table III.8:</b> Result of antibacterial activity of Cam extract with the method of MIC     | 65 |
| <b>Table III.9:</b> Result of antibacterial activity of Cur extract with the method of MIC ... | 65 |
| <b>Table III.10:</b> % inhibition of denaturation of extracted plants .....                    | 68 |
| <b>Table III.11:</b> The growth of hair over the 5 weeks for the control Batch.....            | 70 |
| <b>Table III.12:</b> The growth for hair over the 5 weeks for the treatment Batches.....       | 71 |

# List of figures

## *Bibliography part*

|  |    |
|--|----|
| <b>Figure I.1:</b> Thymelaea hirsuta (Original Photo) .....                          | 6  |
| <b>Figure I.2:</b> Geographical distribution of Thymelaea .....                      | 7  |
| <b>Figure I.3:</b> Biological proprieties of Methane .....                           | 8  |
| <b>Figure I.4:</b> Matrecaria Chamomile (Personal picture).....                      | 9  |
| <b>Figure I.5:</b> Different plant parts and habit of M. Chamomile.....              | 10 |
| <b>Figure I.6:</b> Accepted species of Genus Matricaria and their distribution ..... | 11 |
| <b>Figure I.7:</b> Biologicals proprieties of Chamomile .....                        | 13 |
| <b>Figure I.8:</b> The different part of Curcumin .....                              | 15 |
| <b>Figure I.9:</b> Some therapeutic application of Turmeric .....                    | 16 |
| <b>Figure I.10:</b> Semi-developed structures of curcuminoids.....                   | 17 |

## *Experimental part*

|  |    |
|--|----|
| <b>Figure II.1:</b> Presentation of the studied plants.....                                    | 19 |
| <b>Figure II.2:</b> The washing of the thymelaea.....  | 19 |
| <b>Figure II.3:</b> Vacuum drying chamber .....  | 20 |
| <b>Figure II.4:</b> The used grinder.....  | 20 |
| <b>Figure II.5:</b> The obtained powders .....   | 20 |
| <b>Figure II.6:</b> The hydrodistillation experience.....                                      | 21 |
| <b>Figure II.7:</b> The maceration with ethanol experience.....                                | 21 |
| <b>Figure II.8:</b> The maceration with distilled water experience.....                        | 22 |
| <b>Figure II.9:</b> Curcumin Extraction .....  | 22 |
| <b>Figure II.10:</b> The extracted solutions .....   | 23 |
| <b>Figure II.11:</b> The extracts of plant.....  | 23 |
| <b>Figure II.12:</b> Total polyphenol assay steps.....   | 26 |
| <b>Figure II.13:</b> Total Flavonoids assay steps .....  | 27 |
| <b>Figure II.14:</b> The reaction of vanillin with the terminal flavonoid group of catechins . | 27 |
| <b>Figure II.15:</b> The steps for the determination of condensed tannins .....                | 28 |
| <b>Figure II.16:</b> Mechanical blender .....  | 30 |
| <b>Figure II.17:</b> Adopted cream .....   | 30 |
| <b>Figure II.18:</b> Franz cell.....   | 31 |
| <b>Figure II.19:</b> system for drug release.....  | 31 |
| <b>Figure II.20:</b> Spectrum of absorption of the aqua extract of Thymelaea.....              | 33 |

## ***List of figures***

|  |    |
|--|----|
| <b>Figure II.21:</b> The calibration line of .....                           | 34 |
| <b>Figure II.22:</b> Bacterial strains .....                                 | 36 |
| <b>Figure II.23:</b> Mueller Hinton Medium .....                             | 36 |
| <b>Figure II.24:</b> Preparation of the petri dishes .....                   | 37 |
| <b>Figure II.25:</b> Methods of product's statement .....                    | 37 |
| <b>Figure II.26:</b> Bacterial dishes after incubation .....                 | 38 |
| <b>Figure II.27:</b> Microplates for the MCI method.....                     | 39 |
| <b>Figure II.28:</b> Protocol of the MCI determination .....                 | 40 |
| <b>Figure II.29:</b> Structure of the DPPH radical and its reduced form..... | 41 |
| <b>Figure II.30:</b> Antioxidant activity protocol .....                     | 42 |
| <b>Figure II.31:</b> Presentation of rat's group .....                       | 43 |
| <b>Figure II.32:</b> Injection to induce edema .....                         | 44 |
| <b>Figure II.33:</b> Apparition of edema and their measurement .....         | 44 |
| <b>Figure II.34:</b> Hair removal zone .....                                 | 46 |

### ***Results and discussion part***

|  |    |
|--|----|
| <b>Figure III.1:</b> Extraction yields. ....                         | 48 |
| <b>Figure III.2:</b> Infrared spectrum of thymeleae .....            | 49 |
| <b>Figure III.3:</b> Infrared spectrum of chamomile.....             | 50 |
| <b>Figure III.4:</b> Infrared spectrum of Curcuma.....               | 50 |
| <b>Figure III.5:</b> Total polyphenol content of extracts .....      | 53 |
| <b>Figure III.6:</b> Flavonoid content of various extracts.....      | 54 |
| <b>Figure III.7:</b> Tannin content of different extracts.....       | 54 |
| <b>Figure III.8:</b> Spreading of some creams .....                  | 56 |
| <b>Figure III.9:</b> Liberation curve of Cur from L1 .....           | 57 |
| <b>Figure III.10:</b> Liberation curve of Cur from L2 .....          | 58 |
| <b>Figure III.11:</b> Liberation curve of Cam from L3.....           | 58 |
| <b>Figure III.12:</b> Liberation curve of Cam from L4.....           | 59 |
| <b>Figure III.13:</b> Liberation curve of active agent from L5 ..... | 59 |
| <b>Figure III.14:</b> Liberation curve of active agent from L6.....  | 60 |
| <b>Figure III.15:</b> Liberation curve of Thy from L7.....           | 60 |

## List of figures

|  |    |
|--|----|
| <b>Figure III.16:</b> Results of antibacterial activity against different strains (a, b, c, d: extracted plants, e: L5 and L6, f: L3 and L4..... | 61 |
| <b>Figure III.17:</b> Inhibition diameters of L1.0, L2.0, L1, L2, and the cur extract against the studied strains.....                           | 62 |
| <b>Figure III.18:</b> Inhibition diameters of L3.0, L4.0, L3, L4, and the cam extract against the studied strains.....                           | 63 |
| <b>Figure III.19:</b> Inhibition diameters of L5.0, L6.0, L5, L6, and the cam, cur extracts against the studied strains .....                    | 63 |
| <b>Figure III.20:</b> Inhibition diameters of L7.0, L7, and the thy extract against the studied strains .....                                    | 64 |
| <b>Figure III.21:</b> MIC results for Thymelaea extract .....  | 66 |
| <b>Figure III.22:</b> MIC results for chamomile extract .....  | 66 |
| <b>Figure III.23:</b> Inhibition percentage of free radical DPPH in function of concentration of Cam extract .....                               | 67 |
| <b>Figure III.24:</b> Inhibition percentage of free radical DPPH in function of concentration of Thy extract .....                               | 67 |
| <b>Figure III.25:</b> Paw of rat before treatment .....  | 68 |
| <b>Figure III.26:</b> Rat paw after treatment .....  | 68 |
| <b>Figure III.27:</b> Evolution of the patten diameters .....  | 69 |

## *General Introduction*

### *General Introduction*

Plants have been used for medical purposes for almost 60,000 years. There is evidence that medicinal plants have been cultivated for at least 2,500 years in Greece and Central Asia, and for about 5,000 years in Egypt, China, and India. The use of plants as medicine was first empirical but has progressively developed into a science supported by data. The earliest known written records are from about 5,000 years ago [1].

Despite their well-known therapeutic benefits, medicinal plants can contain hundreds or even thousands of active compounds that are frequently difficult, if not impossible, to detect. While certain ingredients have specific purposes, others have broader applications [2].

Herbal medicine has many advantages over contemporary medicine, reflecting the past's reliance on plants for healing. As antibiotics become less effective against emerging bacteria and viruses, plant-based remedies are becoming more and more popular. As worries about drug side effects increase, herbal medicine is seeing a resurgence. This is especially true when it comes to managing chronic diseases. Ten to twenty percent of hospital admissions are related to adverse reactions to chemical medications [3-5].

The aim of this work is to demonstrate the efficacy of some plants *Thymelaea Hirsuta*, *Matrecaria Chamomile* and *Turmeric* in the pharmaceutical domain, the essential oils of these plants were sought after for this purpose, however no extraction has taken place for these products. To achieve our goal, water-based extracts have been prepared, formulated to galenic forms and tested to evaluate their effectiveness.

This memory will be summarized in three parts:


- The first chapter presents a bibliographic study on medicinal plants and their composition.
- The second chapter describes the methodology used in the extraction, characterization of extracted products, preparation of galenic forms and effectiveness's tests.
- The obtained results and their interpretation are presented in the third chapter.

A global summary succinct the main findings is presented to close.




## *Introduction references*

1. Jamshidi-Kia F., Lorigooini Z., Amini-Khoei, H., Journal of herbmed pharmacology, 2017, 7(1), 1-7.
2. Iserin, P., « Larousse encyclopédie des plantes médicinales », 2001, 2nd edition, Dorling Kindersley Limited, Londres, 241, (11-14).
3. Cosentino S. C. I. G., Tuberoso C. I. G., Pisano B., Satta M. L., Mascia V., Arzedi E., Palmas F., Letters in applied microbiology, 1999, 29(2), 130-135.
4. Danish P., Ali Q., Hafeez M. M., Malik A., Biological and Clinical Sciences Research Journal, 2020, 1(4) 348-355.
5. Cornara L., Borghesi B. A. R., Caporali E., Casazza G., Roccotiello E., Troiano G., Minuto, L., Plant systematics and evolution, 2005, 250, 157-172.



*Chapter I*  
*Bibliography*  
*part*



### **I.1. INTRODUCTION:**

Over the last years, there has been a noticeable increase in fascination with natural plants that have long been used as cures for a variety of illnesses [1].

These days, there is a resurgence of interest in using plant-based remedies. With the help of new experiments and analytical methods in scientific research, the medical field is progressively realizing the effectiveness of traditional prescriptions involving medicinal plants. Ethnobotany, a scientific field that translates common knowledge into scientific understanding, plays a key role in studying traditional herbal medicine [2].

### **I.2. DEFINITION OF MEDICINAL PLANTS:**

The medicinal plant refers to a plant with therapeutic qualities that can be used whole or in plant parts. Some plants may also have used as food, condiment or cosmetic. Medicinal plants used in traditional medicine mainly contain active ingredients for the whole plant, a part of the plant or a mixture of several parts [3]. They are used throughout the world to support, promote, maintain and restore human health. Traditional medicines are part of the wider field of traditional medicine, which includes procedures and practitioners, as well as products [4].

The World Health Organization estimates that 80% of people living in developing countries depend on traditional medicinal practices for their primary health care [5].

### **I.3. THE BENEFITS OF MEDICINAL PLANTS:**

Despite modern medicine's strides, herbal medicine offers numerous benefits, echoing historical reliance on plants for treatment. Today, a resurgence in plant-based treatments emerges as antibiotic efficacy wanes against evolving bacteria and viruses. Herbal medicine sees a revival, notably in chronic disease management, as concerns over medication side effects grow. Approximately 10 to 20% of hospital admissions stem from chemical medication side effects [6].

Interest in medicinal plants has been commonplace since time immemorial. Numerous studies have been carried out in this field by renowned researchers in traditional medicine, some of which are particularly noteworthy [6].

Youdim and Deans examined the effects of aging on antioxidant enzyme activity and fatty acid composition in rat brains, as well as the effects of dietary supplementation with thyme oil or thymol. Untreated rats showed a significant decline in antioxidant enzyme activity with age, while those fed thyme oil or thymol maintained higher levels of antioxidant activity. In addition, rats fed these supplements had higher levels of beneficial fatty acids in their brains, suggesting the beneficial potential of thyme oil as a dietary antioxidant [7].

Cosentino *et al* analyzed the essential oils of different wild thyme species and their antimicrobial activity. Oil components were identified by GC/MS, revealing variable concentrations of hydrocarbon monoterpenes and phenolic monoterpenes. Tests showed antimicrobial activity similar to that of other thyme oils, attributed to their high phenol content. Carvacrol and thymol were the most effective against bacteria, suggesting their potential use in prolonging food shelf life [8].

Lavender's impact on stress levels was investigated by Ghavami *et al*, analyzing 21 studies with a total of 791 intervention group participants and 804 controls. The findings demonstrated a notable decrease in stress scores among those who used lavender compared to controls. Subgroup analyses underscored the efficacy of *L. angustifolia*, student groups, and the perceived stress scale. Overall, the study supports lavender's significant role in stress reduction, advocating for its inclusion in stress management interventions, especially for students [9]. Another research paper proved that the lavender is a rich source of phenolic acids and flavonoids, thus resulting in antioxidant and anti-inflammatory properties [10].

The medicinal properties of Aloe Vera were explored by Danish *et al*, this plant is rich in beneficial compounds. Alcohol extracts from leaves and roots are tested against various bacterial and fungal strains, their results showed promising antibacterial and antifungal activity [11].

#### **I.4. PRINCIPAL INGREDIENTS OF MEDICINAL PLANTS:**

Medicinal plants contain hundreds if not thousands of active chemicals, which are often difficult to identify, even though their medicinal effects are well known. Some of these active ingredients are listed in the bullet points below [12]:

- **Phenols:** diverse compounds in plants, from salicylic acid to phenolic glucoside acids, exhibit anti-inflammatory and antiseptic properties, serving as plant defenses. Phenolic acids like rosmarinic acid offer antioxidant benefits and may combat viruses. Certain plants yield salicylic derivatives and methyl salicylate through distillation, with therapeutic potential.
- **Flavonoids:** pigments found in most plants, have a wide range of beneficial effects. They act as antioxidants, improving blood circulation, and possess anti-inflammatory, antiviral and hepatoprotective properties. Flavonoids such as hesperidin and rutin strengthen capillary walls, while isoflavones, found in plants such as red clover, offer estrogenic effects useful in the treatment of menopausal disorders.
- **Tannins:** present in all plants at different levels, give a bitter taste to bark or leaves, making them unfit for consumption by certain animals. These compounds contract tissues by binding proteins, and are used in tanning hides. They stop bleeding, fight infection and promote healing of damaged tissue, as in the case of eczema or burns. Plants rich in tannins, such as oak and acacia, are used to strengthen supple tissues and drain excessive secretions, as in cases of diarrhea.
- **Essential oils:** obtained by distilling plants, are oxygenated aromatic compounds, sometimes of terpenoid origin. They are widely used in perfumery and possess a variety of properties, such as the antiseptic effect of tea tree.

#### **I.5. THE STUDIED PLANTS:**

The use of medicinal plants remains largely unexploited in Algeria in the cosmetics, pharmaceutical and food sectors [14]. As part of our efforts to enhance the value of Algerian flora, the three plants listed below pique our curiosity.



*I.5.1. Thymelaea hirsuta*


The Thymelaeaceae family, part of the Magnoliopsida includes some 1,200 species in 67 genera. Etymologically, the generic name "Thymelaea" comes from a species whose leaves are reminiscent of thyme and whose fruit is reminiscent of a small "olive", derived from the Greek "elaia".

The Thymelaeaceae are represented by around ten species belonging to the genera Thymelaea and Daphne. They are found in tropical and temperate zones around the world, mainly in Africa, but are absent from colder climates.

In Mediterranean regions, the Thymelaea genus comprises around 31 species of evergreen shrubs and herbs of these, Thymelaea hirsuta is considered the most representative species on the Mediterranean coast [14, 15].

Thymelaea hirsuta Endel was described as a perennial shrub that can reach heights of 2 to 3 meters [16]. Its leaves are very small, densely imbricated, leathery, elongated ovals with sharp points. Like the stems, they are smooth on the underside and pubescent to woolly on top. The flowers, 2 to 5 in number, are found at the tip of the twigs and are characterized by a yellowish calyx that falls rapidly. Flowers can be either unisexual or polygamous, and are borne on different plants.

Thymelaea hirsuta Endel is a hermaphrodite plant. Its fruits are smooth berries that are eaten by animals, ensuring their dispersal (zoochore dispersal). The plant flowers from October to April are pollinated by insects [17]. Two types of flowers can be distinguished, male and female greenish male flowers have a cylindrical calyx, while female flowers are yellowish with a swollen calyx.




**Figure I.1: a: *Thymelaea hirsuta* (Original Photo), b: female flowers, c: male flowers**

The use of thymelaea for hair care is not new and has been passed down through generations in some cultures. Many people, including our grandmothers, have used thymelaea-infused preparations to wash and treat their hair, noticing its beneficial effects as thickening moisturizing and hair growth.

The dried flowering tops of the plant are traditionally ground and then boiled in water to obtain a viscous solution which is spread daily over the skin for treatment of cutaneous infections and noninfectious skin disorders such as eczema. It is also reported as having leishmanicidal and vermifuge properties

Badawy *et al* give The systematic classification of *Thymelaea hirsute* reported in below table and the geographical distribution of *T. hirsuta* grows in the Mediterranean coastal plains. *Thymelaea.hirsuta* is situated from Spain to Greece and Turkey, southern side from Morocco to Egypt and it is also founded in Lebanon and Palestine [19].



**Figure I.2:** Geographical distribution of Thymelaea.

**Table I.1:** the classification of Thymelaea hirsuta.

|                    |               |
|--------------------|---------------|
| <b>Division</b>    | Tracheophyta  |
| <b>Class</b>       | Magnoliopsida |
| <b>Superorder</b>  | Rosanae       |
| <b>Order</b>       | Malvales      |
| <b>Family</b>      | Thymelaeaceae |
| <b>Genus</b>       | Thymelaea     |
| <b>Species</b>     | Hirsuta       |
| <b>Arabic name</b> | Methnane      |

The main components of plants of the Thymelaeaceae family are daphnane diterpenes and coumarins, as well as lignans and phenolics. Daphnane diterpenes are the major biologically active constituents of Thymelaeaceae [5].

Preliminary phytochemical screening of Thymelaea hirsuta leaves, flowers and stems revealed the presence of tannins, alkaloids, steroids, saponins, coumarins, reducing compounds and anthraquinones [18]. so that the contents of phenolics and flavonoids in Thymelaea were

determined in high levels by Djermame on several extracts, including an aqueous extract and a methanol extract [20].

Many studies have proven several biological properties characterized by the methane plant, according to the study conducted by Dehimi [21]. It has been shown to possess anti-cancer, anti-inflammatory, and anti-diabetic properties, it has been also used in folk medicine for its antimelanogenesis, antioxidant, hypoglycemic and antiarthritic [22] and antitumor characteristics [5].

According to a study by Toumi *et al.*, thymelaea has demonstrated promising benefits in preventing cell growth and invasion as well as triggering cell cycle arrest in colorectal cancer cells [23]. The hepatoprotective qualities were also identified [24].



**Figure I.3:** Biological proprieties of Methnane.

### ***1.5.2. Chamomile***

Chamomile is a plant that belongs to the Asteraceae family and can be either annual or perennial, according to Dai *et al.* It is indigenous to Europe and Asia's temperate zones. It is grown all over the world for its excellent nutritional, therapeutic, and cosmetic value. It was in use in ancient Greece, Rome, and Egypt for thousands of years. The extensive usage of this herb in Uyghur medicine was first documented in China [25].

Chamomile has a sweet, herbaceous and slightly fruity aroma, daisy-like flowers with a yellow center 1-1.5 cm long and 12-20 white petals, as described by Bayati *et al* [26]. According to Probadh *et al*, it is a plant that grows upright to a height of 50 to 90 cm and is distributed throughout the world. Its blossoms are highly therapeutic [27].



**Figure I.4:** Matrecaria Chamomile (Personal picture).

Traditionally, chamomile has been used for centuries as an anti-inflammatory, antioxidant, mild astringent and healing medicine. It is also used to treat wounds, ulcers, eczema, gout, skin irritations, bruises, burns, canker sores, neuralgia, sciatica, rheumatic pain, hemorrhoids, mastitis and other ailments. Externally, chamomile has been used to treat diaper rash, cracked nipples, chicken pox, ear and eye infections, disorders of the eyes including blocked tear ducts,




conjunctivitis, nasal inflammation and poison ivy. Chamomile is widely used to treat inflammations of the skin and mucous membranes, and for various bacterial infections of the skin, oral cavity, gums, and respiratory tract [28]. Different parts and behaviors of chamomile were described in the following figure [29].



**Figure I.5:** Different plant parts and habit of *M. Chamomile* **a:** Flowering plant, **b:** Flower head, **c:** Ray floret, **d:** Disk floret, **e:** Capitulum I, **f:** Teeth/petals of disc floret, **g:** Anthers, **h:** Stigma, **i:** Seed, **j:** Fresh flowers.

The chamomile genus consists of 5 species that are primarily found across Europe, Northern Africa, Macaronesia, western, south-western and central Asia, and western North America. These species are known to thrive in various environments such as disturbed land, grassland, regions adjacent to roads and railways, as well as in waste areas and unoccupied spaces [29].



**Figure I.6:** Accepted species of Genus *Matricaria* and their distribution.

**Table I.2:** The scientific classification of Chamomile

|                       |   |
|-----------------------|---|
| <b>Kingdom</b>        | <b>Plantae</b>  |
| <b>Subkingdom</b>     | Tracheobionta   |
| <b>Super division</b> | Spermatophyta   |
| <b>Division</b>       | Magnoliophyta   |
| <b>Class</b>          | Magnoliopsida   |
| <b>Order</b>          | Asterales   |
| <b>Family</b>         | Asteraceae  |
| <b>Genus</b>          | <i>Matricaria</i>   |
| <b>Species</b>        | <i>Chamomilla</i>   |
| <b>Synonyms</b>       | <i>Matricaria recutita</i> L ;<br><i>Chamomilla vulgaris</i> Gray |

The chamomile harbors a diverse range of therapeutically valuable compounds, among these, sesquiterpenes, flavonoids, coumarins, and polyacetylenes stand out as crucial constituents. Noteworthy coumarins in *M. chamomilla* include herniarin, umbelliferone, and other minor variations.

The glucoside precursor of herniarin, (Z)- and (E)-2- $\beta$ -d-glucopyranosyloxy-4-methoxycinnamic acid has been identified as a native compound in chamomile. Additionally, eleven bioactive phenolic compounds, including herniarin, umbelliferone, chlorogenic acid, caffeic acid, apigenin, apigenin-7-O-glucoside, luteolin, luteolin-7-O-glucoside, quercetin, rutin, and naringenin, are present in chamomile extract [30].

The obtained results proved by Stanojevic *et al* the presence of 52 components, wherein the highest content of  $\beta$ -farnesene (29.8 %),  $\alpha$ -farnesene (9.3 %),  $\alpha$ -bisabolol and its oxide (15.7 %), chamazulene (6.4 %), germacrene D (6.2 %) and spiroether (5.6 %) is determined [31].


The phytochemistry composition of essential oils and extracts of *M. chamomilla* has been widely analyzed, showing that the plant contains over 120 constituents. Essential oils are generally composed of terpenoids, such as  $\alpha$ -bisabolol and its oxides A and B, bisabolone oxide A, chamazulene, and  $\beta$ -farnesene, among other compounds. On the other hand, *M. chamomilla* extracts were dominated by phenolic compounds, including phenolic acids, flavonoids, and coumarins [32].

The anti-inflammatory and antioxidant activities of chamomile within the analgesic action are well known, so that this medicinal plant is usually used as liquid extracts, tablets, or capsules to decrease sleep and digestive tract disorders and to diminish anxiety and neuralgic pains (headache, sciatica, rheumatic pain). Topical application is useful in treatment of epidermis and mucous membrane inflammation (burns, sunburn, insect bites, psoriasis, eczema, and conjunctivitis). In addition to the aforementioned properties, recent studies have also shown a possible role of chamomile in the prevention of tumors, cardiovascular diseases, and diabetes [33].

Chamomile tea finds widespread use across Europe, South America, and Mexico, particularly among parents who rely on it to alleviate colic, digestive issues, fever, insomnia, and the discomfort associated with teething in children.

Additionally, women have historically turned to chamomile to address symptoms of menstrual cramps. Its benefits extend beyond ingestion. Chamomile is also valued for its calming properties, often recommended for muscle cramps, headaches, and soothing indigestion and colic-related discomfort.

Inhaling steam infused with chamomile is a common remedy for respiratory tract irritations. Moreover, chamomile-based creams and ointments are prized for their ability to soothe and heal various skin ailments, including burns, wounds, diaper rash, and sore nipples. Even in homeopathic form, such as tablets, chamomile is administered to babies to ease teething discomfort and fussiness [34].



**Figure I.7:** Biologicals proprieties of Chamomile.

### 1.5.3. *Curcuma longa*

Turmeric was first utilized approximately 4,000 years ago in India's Vedic culture, when it was both a culinary spice and a holy herb. It most likely made its way to East Africa, West Africa, China, and Jamaica by the eighteenth century. When Marco Polo first wrote about this spice in

1280, he was amazed at how closely the vegetable resembled saffron in its properties. Turmeric has a long history of medicinal use in South Asia, according to Ayurvedic and Unani traditions, as well as Sanskrit medical treatises [35].

A popular spice that has captured the attention of both the medical and culinary communities. Turmeric, a perennial herbaceous plant with rhizomes (*Curcuma longa*) belonging to the ginger family, has been recognized for its healing properties for centuries. While the medicinal benefits of turmeric and its active component, curcumin, have been understood for ages, scientists have only recently begun exploring its precise mechanisms of action and identifying its bioactive elements [36].

Turmeric, known as haldi is a vibrant yellow-orange spice derived from the rhizome of the *Curcuma longa* plant. This plant typically grows up to 3 feet tall and features lance-shaped leaves. It produces spikes of yellow flowers that emerge from a fleshy rhizome or an underground stem. The medicinal powder of turmeric comes from the orange pulp found within the rhizome [37]. In the next, the content of turmeric is described [39]:

- **The Rhizome:** *Curcuma longa* has numerous underground stems, including a main one measuring around 5 cm x 2.5 cm. At maturity, it produces many aromatic, cylindrical, or ellipsoidal lateral rhizomes, with a gray-yellow exterior and a dark orange interior.
- **The Leaf:** The leaves are typically wide (7 to 25 cm) and long (up to 50 cm), uniformly green in color. They are arranged in two rows, alternating with each other. Their spikes can reach up to 20 cm, and the flowers are sterile



**Figure I.8:** The different part of Curcumin.

In traditional medicine, turmeric has been used in therapeutic preparations over the centuries in different parts of the world. In Ayurvedic practices, turmeric is thought to have many medicinal properties including strengthening the overall energy of the body, relieving gas, dispelling worms, improving digestion, regulating menstruation, dissolving gallstones, and relieving arthritis [39].

The composition of turmeric encompasses a diverse array of components, with over 100 identified thus far. The primary constituents include a volatile oil rich in turmerone, alongside curcuminoids, such as curcumin, demethoxycurcumin, 5'-methoxycurcumin, and dihydrocurcumin, which exhibit natural antioxidant properties. Standard turmeric comprises moisture (>9%), curcumin (5–6.6%), minimal extraneous matter (<0.5% by weight), low levels of mold (<3%), and volatile oils (<3.5%), containing various compounds like d- $\alpha$ -phellandrene, d-sabinene, cinol, borneol, zingiberene, and sesquiterpenes. These sesquiterpenes, including germacrone, termerone,  $\alpha$ -, and  $\beta$ -termerones, among others, often exhibit species-specific characteristics [40].





**Figure I.9:** Some therapeutic application of Turmeric.

The aromatic essence of turmeric primarily derives from turmerone, arturmerone, and zingiberene. Additionally, the rhizomes contain polysaccharides-ukonans, stigmasterole,  $\beta$ -sitosterole, cholesterol, and 2-hydroxymethyl anthraquinone. In terms of nutrition, 100 grams of turmeric provides 390 kcal, 10 grams of total fat, 3 grams of saturated fat, essential minerals like calcium, phosphorus, sodium, potassium, iron, and vitamins such as thiamine, riboflavin, niacin, and ascorbic acid. Furthermore, turmeric is recognized as a notable source of the  $\omega$ -3 fatty acid and  $\alpha$ -linolenic acid [40].

The main component found in turmeric is called curcumin, which is known for its diverse range of healing properties [41]. Curcumin presents different therapeutic action as well as hepatoprotective [42], antimicrobial and cardio protective [43], anticancer [44], hypoglycemic [45] and antiarthritic properties [46].

Jovanovic *et al* have determined that turmeric is a donor of proton of the central methyl group rather than phenolic groups in aqueous, acidic and acetonitrile solutions, which explains the potent antioxidant activity of curcumin, as demonstrated by its attribution to the neutralization of




free radical [47]. However, Barclay *et al* proposed that curcumin's antioxidant properties stem from phenolic groups that have ability to give a proton [48].

According to Priyadarsin *et al*, the presence of methoxy groups enhances this property even further and the phenolic groups play a crucial role in the curcumin's antioxidant activity [49].

Another research by Mou-Tuan Huang *et al* has demonstrated that curcumin inhibits tumor growth, protease C activity, cyclooxygenase epidermal in vitro activity and epidermal inflammation [50]. Additionally, curcumin exhibits tremendous promise in the treatment of Alzheimer's disease [51].

Since the Covid-19 pandemic struck Wuhan, numerous research projects have been launched around the world and the fight against the virus has been the focus of extensive studies. One of them, is being conducted by Khaerunnisa team [52], which evaluated the bioactive components found in medicinal plants as curcumin, that may be Covid-19 inhibitors using a molecular modeling technique. They found that, when combined with other natural compounds, curcumin appears to have the most potential for acting as Covid-19 inhibitors.



**Figure I.10:** Semi-developed structures of curcuminoids.

## *Bibliographical references*

1. ABBOU FAIZA, A. R. (06octobre2018). Evaluation de l'activité antioxydante et antidiabétique des extraits de la partie aérienne de *Mentha pulegium* L. *Journée nationale « la recherche et la valorisation des plantes d'intérêts écologique. Economique et patrimonial*», 19.
2. LAHSISSENE H., K. A. (décembre 2009). catalogue des plantes médicinales utilisées dans la région de zaer. *revue de Botanique ( en ligne)*, 186.
3. Sucher, N. J., & Carles, M. C. (2008). Genome-based approaches to the authentication of medicinal plants. *Planta medica*, 74(06), 603-623 Gazengel, J. M., & Orecchioni, A. M. (2013). *Le préparateur en pharmacie*. Technique et Documentation
4. Organization, World Health. 2015. WHO traditional medicine strategy 2002–2005. Geneva: World Health Organization; 2002
5. Sanna, G., Madeddu, S., Murgia, G., Serreli, G., Begala, M., Caboni, P., ... & Giliberti, G. (2020). Potent and selective activity against human immunodeficiency virus 1 (HIV-1) of *Thymelaea hirsuta* Extracts. *Viruses*, 12(6),664.
6. Iserin, P. (2001). Larousse encyclopédie des plantes médicinales. *Identification, Préparations, soins. 2nd edition, Dorling Kindersiey Limited, Londres*, 241
7. Youdim, K. A., & Deans, S. G. (2000). Effect of thyme oil and thymol dietary supplementation on the antioxidant status and fatty acid composition of the ageing rat brain. *British Journal of Nutrition*, 83(1), 87-93.
8. Cosentino, S. C. I. G., Tuberoso, C. I. G., Pisano, B., Satta, M. L., Mascia, V., Arzedi, E., & Palmas, F. (1999). In-vitro antimicrobial activity and chemical composition of Sardinian thymus essential oils. *Letters in applied microbiology*, 29(2), 130-135.
9. Ghavami, T., Kazemini, M., & Rajati, F. (2022). The effect of lavender on stress in individuals: A systematic review and meta-analysis. *Complementary therapies in medicine*, 68, 102832.
10. Dobros, N., Zawada, K. D., & Paradowska, K. (2022). Phytochemical profiling, antioxidant and anti-inflammatory activity of plants belonging to the *Lavandula* genus. *Molecules*, 28(1), 256.

## ***Bibliographical references***

11. Danish, P., Ali, Q., Hafeez, M. M., & Malik, A. (2020). Antifungal and antibacterial activity of aloe vera plant extract. *Biological and Clinical Sciences Research Journal*, 2020(1).
12. Iserin, P. (2001). Larousse encyclopédie des plantes médicinales. *Identification, Préparations, soins. 2nd edition, Dorling Kindersiey Limited, Londres*, 241. (11-14).
13. Miara, M. ., (2013). Phytothérapie et taxonomie des plantes médicinales spontanées dans la région de Tiaret (Algérie). *Phytothérapie*, 206-218.
14. Borris, R. B. (1988). Ethnopharmacologic and phytochemical studies of the Thymelaea . *Ethnopharmacology* , 41-49.
15. Herbada, D. (2006). Origin and diversification of Thymelaea, inferences from a phylogenetic study based on ITS(rDNA)sequences. *plant systematics and evolution* , 159-187.
16. BOUNAB SOUHILA, L. T. (octobre 2019). Chemical composition and antibacterial activity of essential oil of Thymelaea hirsuta from Algetia. *BIODIVERITAS* , 2868-2876.
17. Cornara, L., Borghesi, B. A. R. B. A. R. A., Caporali, E., Casazza, G., Roccotiello, E., Troiano, G., & Minuto, L. (2005). Floral features and reproductive ecology in Thymelaea hirsuta (L.) Endl. *Plant systematics and evolution*, 250, 157-172.
18. Amari, N. O., Razafimandimby, B., Auberon, F., Azoulay, S., Fernandez, X., Berkani, A., ... & Landreau, A. (2021). Antifungal and antiaging evaluation of aerial part extracts of Thymelaea hirsuta (L.) endl. *Natural Product Communications*, 16(2), 1934578X20987932.
19. Ahmed M Badawy, H. A. (2019). Review article on chemical constituents and biological activity of Thymelaea hirsuta. *RECORDS OF PHARMACEUTICAL AND BIOMEDICAL SCIENCES* , 28-32.
20. Djermane, N. ., (2020). Chemical composition and in vitro evaluation of antioxidant, antimicrobial, and enzyme inhibitory activities of Erucaria uncata and Thymelaea hirsuta. *Biocatalysis and Agricultural Biotechnology*.

## *Bibliographical references*

21. DEHIMI, K. (2018). Study of biological activities of extracts from *Thymelaea microphylla* Coss. and Dur (Doctoral dissertation),
22. Anti-microbial screening and cytotoxic activity of aerial part of *Thymelaea hirsuta* L. essential oil growing in south-west Tunisia. Samir Felhi<sup>1</sup>, Mouna Chaaibia<sup>1</sup>, Sana Bakari<sup>1</sup>, Riadh Ben Mansour<sup>2</sup>, Ahmed Békir<sup>3</sup>, Néji Gharsallah<sup>1</sup> and Adel Kadri<sup>1,4\*</sup>2017
23. Toumi, I., Yatouji, S., Borie, N., Remy, S., Renault, J. H., Chazee, L., ... & El Btaouri, H. (2023). Anti-tumoral Effect of *Thymelaea hirsuta* L. Extracts in Colorectal Cancer Cells. *Anti-Cancer Agents in Medicinal Chemistry (Formerly Current Medicinal Chemistry-Anti-Cancer Agents)*, 23(6), 687-698
24. Azza, Z., Marnissi, F., Naya, A., Benjelloun, N., Zamyati, S., Amrani, M., & Oudghiri, M. (2012), Toxicological evaluation of *Thymelaea hirsuta* and protective effect against CCl<sub>4</sub>-induced hepatic injury in rats. *International Journal of Biological and Chemical Sciences*, 6(1), 379-393.
25. Dai, Y. L., Li, Y., Wang, Q., Niu, F. J., Li, K. W., Wang, Y. Y., ... & Gao, L. N. (2022). Chamomile: a review of its traditional uses, chemical constituents, pharmacological activities and quality control studies. *Molecules*, 28(1), 133
26. Jalal Bayati Zadah and Nasroallah Moradi KOr, Z. M. (2014). Chamomile ( *MAtricaria recutita*) As a Valuable Medicinal Plant. *INternational Journal of Advanced Biological and Biomedical Research*, 823-829.
27. Prabodh Satyal, S. S. (2015). Composition and Bioactivities of an (E)- - FArnesene Chemotype of Chamomile (*MAtricaria chamomilla* ) Essential Oil from Nepal. *Natural Product Communication* , 1453-1457
28. JanMeJai K. SrivaSTava, e. S. (August 30, 2010). Chamomile: A herbal medicine of the past with a bright future (Review). *Molecular Medicine rePorTS*, 896.
29. Chauhan, R., Singh, S., Kumar, V., Kumar, A., Kumari, A., Rathore, S., ... & Singh, S. (2021). A comprehensive review on biology, genetic improvement, agro and process technology of German chamomile (*Matricaria chamomilla* L.). *Plants*, 11(1), 29
30. Chamomile (*Matricaria chamomilla* L.): An overview Ompal Singh, Zakia Khanam<sup>1</sup>, Neelam Misra, Manoj Kumar Srivastava 2010.

## *Bibliographical references*

31. Stanojevic, L. P., Marjanovic-Balaban, Z. R., Kalaba, V. D., Stanojevic, J. S., & Cvetkovic, D. J. (2016). Chemical composition, antioxidant and antimicrobial activity of chamomile flowers essential oil (*Matricaria chamomilla* L.). *Journal of essential oil bearing plants*, 19(8), 2017-2028.
32. El Mihaoui, A., Esteves da Silva, J. C., Charfi, S., Candela Castillo, M. E., Lamarti, A., & Arnao, M. B. (2022). Chamomile (*Matricaria chamomilla* L.): a review of ethnomedicinal use, phytochemistry and pharmacological uses. *Life*, 12(4), 479.
33. Catani, M. V., Rinaldi, F., Tullio, V., Gasperi, V., & Savini, I. (2021). Comparative analysis of phenolic composition of six commercially available chamomiles (*Matricaria chamomilla* L.) extracts: Potential biological implications. *International Journal of Molecular Sciences*, 22(19), 10601.
34. (Jalal Bayati Zadah and Nasroallah Moradi KOr, 2014)
35. Yadav, R. P., & Tarun, G. (2017). Versatility of turmeric: A review the golden spice of life. *Journal of Pharmacognosy and Phytochemistry*, 6(1), 41-46.
36. Hewlings, S. J., & Kalman, D. S. (2017). Curcumin: A review of its effects on human health. *Foods*, 6(10), 92.
37. Nagpal, M., & Sood, S. (2013). Role of curcumin in systemic and oral health: An overview. *Journal of natural science, biology, and medicine*, 4(1), 3.
38. Romain Leroy. La curcumine : provenance et potentiels thérapeutiques. Sciences pharmaceutiques.2019. ffdumas-02480721
39. Prasad, S., & Aggarwal, B. B. (2011). Turmeric, the golden spice. *Herbal Medicine: Biomolecular and Clinical Aspects. 2nd edition*.
40. Prasad, S., & Aggarwal, B. B. (2011). Turmeric, the golden spice. *Herbal Medicine: Biomolecular and Clinical Aspects. 2nd edition*.
41. Nagpal, M., & Sood, S. (2013). Role of curcumin in systemic and oral health: An overview. *Journal of natural science, biology, and medicine*, 4(1), 3.
42. Miyakoshi M, Yamaguchi Y, Takagaki R. et al. Hepatoprotective effect of sesquiterpenes in turmeric. *Biofactors*. 2004; 21:167–70
43. Mohanty I, Arya D. S, Gupta S. K. Effect of *Curcuma longa* and *Ocimum sanctum* on myocardial apoptosis in experimentally induced myocardial ischemic-reperfusion injury. *BMC Complement Altern Med*. 2006; 6:3.

## *Bibliographical references*

44. Azuine M. A, Bhide S. V. Adjuvant chemoprevention of experimental cancer: Catechin and dietary turmeric in forestomach and oral cancer models. *J Ethnopharmacol.* 1994; 44:211–7. [[PubMed](#)] [[Ref list](#)]
45. Honda S, Aoki F, Tanaka H. Et al. Effects of ingested turmeric oleoresin on glucose and lipid metabolisms in obese diabetic mice: A DNA microarray study. *J Agric Food Chem.* 2006; 54:9055–62
46. (Funk J. L, Frye J. B, Oyarzo J. N. et al. Efficacy and mechanism of action of turmeric supplements in the treatment of experimental arthritis. *Arthritis Rheum.* 2006; 54:3452–6.
47. S. V. Jovanovic, S. Steenken, C. W. Boone, and M. G. Simic, “H-atom transfer is a preferred antioxidant mechanism of curcumin,” *J. Am. Chem. Soc.*, vol. 121, no. 41, pp. 9677–9681, Oct. 1999.
48. L. R. C. Barclay et al., “On the antioxidant mechanism of curcumin: Classical methods are needed to determine antioxidant mechanism and activity,” *Org. Lett.*, vol. 2, no. 18, pp. 2841–2843, Sep. 2000.
49. K. I. Priyadarsini et al., “Role of phenolic O-H and methylene hydrogen on the free radical reactions and antioxidant activity of curcumin,” *Free Radic. Biol. Med.*, vol. 35, no. 5, pp. 475–484, Sep. 2003.
50. M.-T. Huang, T. Lysz, T. Ferrara, T. F. Abidi, J. D. Laskin, and A. H. Conney<sup>2</sup>, “Inhibitory Effects of Curcumin on in Vitro Lipoxygenase and Cyclooxygenase Activities in Mouse Epidermis<sup>1</sup>,” pp.813-819, 1991.
51. J. Jean-Pierre, “Curcuma et curcumine: de l’histoire aux intérêts thérapeutiques,” *Univ. Caen, Normandie*, pp. 51–55, Dec. 2015.
52. S. Khaerunnisa, H. Kurniawan, R. Awaluddin, and S. Suhartati, “Potential Inhibitor of COVID-19 Main Protease (M pro) from Several Medicinal Plant Compounds by Molecular Docking Study,” *Preprints*, no. March, pp. 1–14, 2020.

*Chapter II*  
*Experimental*  
*part*





## II.1. INTRODUCTION:

Plants have played an important role in medicine since ancient times, offering a vast of therapeutic compounds that have been utilised for healing purposes. These compounds include alkaloids, flavonoids, terpenes and phenolics, among others. Many of these compounds have been isolated and studied for their pharmacological effects. In our work, our plan is to attempt extractions from some plants or work with natural extracts.

## II.2. MATERIALS AND METHODS:

### II.2.1. Chemical products:

**Table II.1:** chemical products.

|                    |                    |
|--------------------|--------------------|
| Almond oil         | salicylic acid     |
| coco betaine       | sodium bicarbonate |
| copper sulfate     | starch             |
| Fenugreek oil      | tri-ethanolamine   |
| kathon CG          | vasline            |
| lavender oil       | vitamin E          |
| magnesium stearate | zinc oxide         |
| propylene glycol   | zinc sulfate       |

### II.2.2. Vegetal materials:

Our study focused on three plants:

- The first known as “wild thymelaea”. The upper part of this plant was harvested in the flowering period from the Rahouia region, specifically Ain Abd Errahim, 40km from state of Tiaret, on December.

- The second is Chamomile, with its delicate flowers and gentle fragrance, it is commonly used in teas and herbal remedies for its calming effect, anti-inflammatory and antioxidant benefits. It's purchased from the market.
- The third is derived from the *Curcuma longa* plant, it's celebrated for its potential health benefits as anti-inflammatory and antioxidant properties. It's also purchased from the market.

**A****B****C**

**Figure II.1:** Presentation of the studied plants A: Thymelaea, B: Chamomile, C: *Curcuma longa*.

### II.2.3. Methodology of work

These studies were carried out at the chemistry laboratory at the faculty of Materials Sciences of Ibn Khaldoun University of Tiaret

- **Washing process:** the washing process of the plant involved repeated washing at least three times to ensure that all impurities are removed from the soil and sediment followed by final wash with distilled water.



**Figure II.2:** The washing of the thymelaea.

- **Drying process:** the plant was dried in an air dryer at a temperature of 40-45°C for 48 hours.



**Figure II.3:** Vacuum drying chamber.

- **Grinding process:** The plants mentioned above were milled in the same manner using the grinder depicted below.



**Figure II.4:** The used grinder.



**Figure II.5:** The obtained powders A: Thymelaea, B: Chamomile, C: Curcuma longa.

➤ *Hydrodistillation experience*

The obtained powders of Thymelaea and Chamomile was subjected to hydrodistillation for three hours. the process was repeated four times and a homogenized filtrate. However, it was impossible to extract the oil.



**Figure II.6:** The hydrodistillation experience.

➤ *Maceration with ethanol*

This process facilitates the extraction of active compounds but with our plants the oils weren't obtained. In this experience, the maceration for 48 hours includes immersing Thymelaea and Chamomile in ethanol maintained at 4°C was studied.



**Figure II.7:** The maceration with ethanol experience.

➤ *Maceration with distilled water*

In this part, the maceration at room temperature for 24 hours involves immersing Thymelaea and Chamomile in distilled water with vigorous agitation was released three times as described in previous paper [1] but the oil wasn't also obtained.



**Figure II.8:** The maceration with distilled water experience.

➤ *Extraction of curcumin*

We placed the obtained curcuma powder with acetone in the reflux set-up for 24 hours at a temperature of 35 °C.



**Figure II.9:** Curcumin Extraction.



Several methods were used to extract essential oils from the studied plants, but no results were obtained. Therefore, we decided to use extract of plant in our work.

#### II.2.4. Preparation of the extract:

After maceration with distilled water, the mixture was filtered then the obtained solution was dried in vacuum drying chamber for 24h at 40°C.



**Figure II.10:** The extracted solutions A: Thymelaea, B: Chamomile, C: Curcuma longa.



**Figure II.11:** The extracts of plant A: Chamomile, B: Thymelaea, C: Curcuma longa.

### II.3. QUALITATIVE ANALYSIS

#### IR analysis of extracts

Infrared (IR) spectroscopy serves as a pivotal analytical method primarily utilized for identifying functional groups in organic molecules and determining the structures of certain simple molecules. This technique relies on the absorption of infrared radiation by the sample under examination, allowing for the detection of characteristic vibrations of chemical bonds. These

vibrations aid in the analysis of the chemical functionalities present within the sample. Leaf extracts from three species were subjected to IR analysis within the range of 400 to 4000  $\text{cm}^{-1}$ , utilizing an ALPHA-P BRUKER SPECTROMETER. The analyses took place at the materials chemistry laboratory of the University of Oran1.

#### II.4. PHYTOCHEMICAL SCREENING

This study highlights the presence of some groups of diverse metabolites in plants and to detect these latter we perform a series of chemical analyses of the obtained extract and among these compounds we mention: alkaloids, phenolic compounds and tannins (reaction with ferric chloride), flavonoids, saponins and coumarins [2-5].

- ✚ Alkaloid test: 2mL of the extracted solution was mixed with 2 mL of concentrated hydrochloric acid (HCl) followed by the addition of a few drops of Mayer's reagent, the formation of a green or white precipitate indicates the presence of alkaloids.
- ✚ Coumarins test: 1 mL of the extracted solution was mixed with 1 mL of sodium hydroxide 10%. The formation of a yellow color indicates the presence of coumarins.
- ✚ Flavonoids test: 1 mL of sodium hydroxide solution 10% was added to 1 mL of the aqueous extract. The presence of flavonoids is indicated by the formation of a yellow to orange color.
- ✚ Phenolic test: 1 mL of extracted solution was diluted with 2 mL of distilled water, followed by the addition of a few drops of a ferric chloride solution 10%. The formation of a blue or green color indicates the presence of phenols.
- ✚ Quinone test: 1 mL of the extracted solution was merged with 1 mL of concentrated sulfuric acid. The presence of quinones is indicated by the formation of a red color.
- ✚ Saponins test: 1 mL of aqueous extract was diluted with 5mL of water, the tube was shaken for 2 minutes. The formation of stable foam after 15 min confirms the presence of saponins.
- ✚ Steroid test: 0,5 mL of the extracted solution was treated with drops of chloroform, acetic anhydride and concentrated sulfuric acid. The formation of a dark pink color indicates the presence of steroids.
- ✚ Tannin test: 2 mL of the extracted solution was diluted with distilled water and 2-3 drops of ferric chloride 5% ( $\text{FeCl}_3$ ) were added. A green-black or blue coloration would indicate the presence of tannin.



## II.5. Quantitative analysis of extracts

Colorimetric methods based on UV-visible spectrophotometry were used to assess the quantity of phenolic compounds: total polyphenols, total flavonoids and tannins present in the leaves of the studied species [6-8].

### II.5.1. Dosage of total polyphenols

The principle of the method: the dosage of polyphenols is released by spectral photometry using the Folin-Ciocalteu method. This method is based on the oxidoreduction reactions with Folin-Ciocalteu acid which has a yellow color, it is composed of a mixture of phosphotungsthenic and phosphomolybdic acids. The phenol reduces this reagent to a mixture of blue oxides of tungsten and molybdenum that absorb at 765 nm. The intensity of color is proportional to the content of oxidized phenolic compounds.

- ✚ *Operating mode:* Polyphenols were determined using a modified Folin-Ciocalteu photometric assay [9]. The assay protocol for total polyphenols is summarized in figure II.12 below. All manipulations are carried out in triplicate. the blank is also prepared under the same conditions, replacing the quantity of extract with distilled water.
- ✚ *Expression of results:* Total polyphenol concentrations are calculated using the equation developed by Nirmala *et al.* [9].


$$\text{TPC (mg/g)} = \frac{C*V}{m}$$

TPC: total polyphenols content.

C: extract concentration established from calibration curve.

V: extract volume

m: mass of dry extract



**Figure II 12:** Total polyphenol assay steps.

## II.5.2. Determination of total flavonoids

- ✚ *Principle of the method:* The determination of total flavonoids is based on a colorimetric test using aluminum trichloride  $\text{AlCl}_3$  as the reagent. Flavonoids have a free hydroxyl (OH) group in position 5, which is able to form a colored complex with aluminum chloride, together with the CO group. Flavonoids form yellow complexes by chelating metals (aluminum). The Al metal loses 2 electrons to join two oxygen atoms of the phenolic molecule acting as an electron donor.
- ✚ *Operating mode:* The total flavonoid content of the extracts was determined using the method described by Marsoul and Kim with a few modifications [10,11]. All manipulations are carried out in triplicate. The blank is also prepared under the same conditions, replacing the quantity of extracted solution with distilled water. The assay protocol for total flavonoid is summarized in the figure below.
- ✚ *Expression of results:* Flavonoid quantification was based on the equation established by Singh *et al.* [12]


$$\text{TFC (mg/g)} = \frac{C \cdot V}{m}$$

TFC: total flavonoid content.

C: extract concentration established from calibration curve.

V: extract volume.


m: mass of dry extract.



**Figure II.13:** Total Flavonoids assay steps.

### II.5.3. Dosage of condensed tannins

✚ *Principle of Method:* Condensed tannins are determined by the vanillin method in acid medium. The principle of this assay is based on the reaction of vanillin with the terminal flavonoid group of catechins, resulting in the formation of a red complex. This is explained by the property of tannins to transform into red anthocyanidols, which absorb light at 500 nm.



**Figure II.14:** The reaction of vanillin with the terminal flavonoid group of catechins.

✚ *Operating mode:* Total condensed tannin content was determined by the Sun at a procedure [13]. All manipulations are carried out in triplicate. the blank is also prepared under the same conditions, replacing the quantity of extract with distilled water.

The assay protocol for total flavonoid is summarized in the figure below.

✚ *Expression of results:* The concentration of condensed tannins is calculated using the equation established by Madhav *et al.* [14].


$$TC \text{ (mg/g)} = \frac{C \cdot V}{m}$$

TC: tannin content (mg ECT/g of dried extract).

C: extract concentration established from calibration curve.

V: extract volume.

m: mass of dry extract.



**Figure II.15:** The steps for the determination of condensed tannins.

## II.6. GALENIC FORMS FORMULATION:

Our work's goal is to create herbal remedies based on plant extracts and assess their therapeutic potential. This section displays the composition of our prepared formulations in each table.

**Table II.2.** The composition of L1.

| Ingredients | Curcuma | Paraffin oil | Vaseline | Magnesium stearate | Starch | CG Kathon |
|-------------|---------|--------------|----------|--------------------|--------|-----------|
| Quantity    | 0.8 g   | 20 mL        | 20 g     | 0.2g               | 4g     | 1 mL      |

**Table II.3.** The composition of L2.

| Ingredients | Curcuma | Paraffin oil | Vaseline | Magnesium stearate | Starch | CG Kathon | Zinc oxide |
|-------------|---------|--------------|----------|--------------------|--------|-----------|------------|
| Quantity    | 0.8 g   | 20 mL        | 20 g     | 0.2g               | 4g     | 1 mL      | 0.8 g      |

**Table II.4.** The composition of L3.

| Ingredients | chamomile | Paraffin oil | Vaseline | Magnesium stearate | Starch | CG Kathon | Flower water |
|-------------|-----------|--------------|----------|--------------------|--------|-----------|--------------|
| Quantity    | 0.8 g     | 15mL         | 22 g     | 0.2g               | 4g     | 1 mL      | 20mL         |

**Table II.5.** The composition of L4.

| Ingredients | chamomile | Paraffin oil | Vaseline | Magnesium stearate | Starch | CG Kathon | Flower water |
|-------------|-----------|--------------|----------|--------------------|--------|-----------|--------------|
| Quantity    | 1 g       | 20mL         | 15 g     | 0.2g               | 5g     | 1.5mL     | 22mL         |

**Table II.6.** The composition of L5.

| Ingredients | Curcuma | chamomile | Vaseline | Magnesium stearate | Starch | CG Kathon | Flower water |
|-------------|---------|-----------|----------|--------------------|--------|-----------|--------------|
| Quantity    | 0.8 g   | 0.7 g     | 23 g     | 0.2g               | 5.5 g  | 1 mL      | 13 mL        |

**Table II.7.** The composition of L6.

| Ingredients | Curcuma | chamomile | paraffin oil | Vaseline | Magnesium stearate | Starch | CG Kathon | Flower water |
|-------------|---------|-----------|--------------|----------|--------------------|--------|-----------|--------------|
| Quantity    | 0.9 g   | 0.8 g     | 8 mL         | 20 g     | 0.2g               | 4 g    | 1 mL      | 13 mL        |

**Table II.8.** The composition of L7.

| Ingredients | Thymelaea | Sodium bicarbonate | copper sulfate | Vaseline | Magnesium stearate | Starch | CG Kathon | Flower water |
|-------------|-----------|--------------------|----------------|----------|--------------------|--------|-----------|--------------|
| Quantity    | 0.9 g     | 0.2 g              | 0.3 mL         | 20 g     | 0.2g               | 8 g    | 1 mL      | 10 mL        |

**Table II.9.** The composition of Shampoo.

| Ingredients | Thymelaea | propylene glycol | coco betaine | CG Kathon | Flower water |
|-------------|-----------|------------------|--------------|-----------|--------------|
| Quantity    | 0.5 g     | 0.2 g            | 6 ml         | 1.5 mL    | 3.5 mL       |

In order to prepare all galenic formulations, we mixed the ingredients in a mechanical blender as follows:



**FigureII.16:** Mechanical blender.



**FigureII.17:** Adopted cream.

## **II.7. QUALITY CONTROL FOR GALENIC FORMS:**

Quality control for galenic forms is necessary to ensure the efficacy, consistency of the prepared formulations. the prepared formulations exhibit the following traits:


- **Consistency and Homogeneity**, Testing the consistency involved placing each product detail on a glass plate.
- **Color and odor** are tested by visual inception
- **Stability** is tested by centrifugation of galenic forms at different times (5min and 30min)
- **pH Level:** all pH galenic forms are determined by pH-meter.

## **II.8. DRUG RELEASE OF GALENIC FORMS:**


### **II.8.1. Operating procedure:**

In order to release the active ingredients from the prepared galenic forms, a system that resembles the Franz cell was attempted. This cell contains a donor compartment, separated from a recipient compartment by the membrane of the skin to be studied. In the donor compartment, the

formulation containing the active ingredient is placed, and successive samples taken from the recipient compartment enable the kinetic profile of release to be monitored over time.



**Figure II.18:** Franz cell.



**Figure II.19:** system for drug release (1. Syringe for sampling, 2. Thermometer, 3. Membrane, 4. pH= 5.5, 5. Stirrer).




The formulations are placed in the donor compartment, while the recipient compartment contains a medium thermostated at 37°C with a pH=5.5 simulated to the skin medium, and magnetic stirring. The system is carefully sealed with paraffin paper and aluminum foil to prevent evaporation of the medium. The active ingredient is thus released by contact of the formulations with the reconstituted medium through the membrane. Over time, 1 mL samples are taken from the recipient compartment using a syringe, then they are diluted in the same medium to a total volume of 10 mL, maintained at a pH of 5.5, and analyzed by UV spectrophotometry. For all experiments, samples are taken over seven hours.

The controlled conditions in all experiments are:

- ***Stirring the medium:*** To ensure uniform concentration of the solution at all points, a magnetic stirrer is used in the flask. The speed of rotation is set at 500 rpm for all experiments. This is crucial, as the absence of stirring would lead to the formation of a concentration gradient in the solution, which could induce errors in UV spectrophotometric analyses.
- ***Temperature:*** Temperature plays an essential role in diffusion phenomena, influencing the solubility of the active ingredient and facilitating diffusion. All experiments are carried out at a constant temperature of 37°C, which corresponds to human body temperature. A heated stirrer is used to keep this temperature
- ***The nature of the medium, its pH and volume:*** Like temperature, the nature of the medium plays a crucial role in the solubility of the active agent, which in turn affects the diffusion process. In addition, the pH of the medium has a significant impact on the hydrolysis rate and solubility of the active ingredient.

### II.8.2. Principe active analysis

- **$\lambda$  max research :**  $\lambda$  max is determined by performing a U.V. spectrum at a concentration of  $10^{-3}$  g/L of each extract in the study medium at pH= 5.5. The UV spectrum of Thymeleae is given in figure II.19, the others are summarized in annex.



**Figure II.20:** Spectrum of absorption of the aqua extract of Thymelaea.

The  $\lambda$  max values thus found ( $\lambda_1$  max = 273 nm /  $\lambda_2$  max = 272nm /  $\lambda_3$  max = 274nm) are correspond respectively to cam, cur and thy. The instrument used is the U.V JENWAY spectrophotometer (Synthesis and Catalysis Laboratory, Ibn Kaldoun University, Tiaret).

- **Calibration line and Calculation of  $\epsilon$  max:** Solutions at known concentrations were analyzed at  $\lambda$  max. The spectral absorption curve of the extract in a medium at pH=5.5 is shown below. Optical density (O.D.) or absorbance (A) is related to concentration (g/L) by the Beer-Lambert law:


$$(\text{D.O})_{\text{max}} = \log (I_{\text{max}}/I_0) = \epsilon_{\text{max}} \cdot l \cdot C$$

$$(\text{D.O})_{\text{max}} = \epsilon_{\text{max}} \cdot l \cdot C$$

- $I_{\text{max}}/I_0$ : is the transmittance.
- $\epsilon$  max: the specific absorption coefficient ( $\text{L}\cdot\text{cm}^{-1} \cdot \text{mole}^{-1}$ ).

- C: concentration of the solution in mole/l.
- l: length of quartz cell (1cm).

The value of  $\epsilon_{\max}$  corresponds to the value of the tangent of the line (D.O)  $\max=f(C)$ . the calibration line of Thymeleae, is cited at **figure II.20** and the other lines are summarized in annex.



**Figure II.21:** The calibration line of Thymeleae.

- **Analysis of transferred quantities**

Release kinetics were studied using a UV-VIS spectrometer (UV 1650PC SCHIMADZU). This device is first calibrated to the wavelength  $\lambda_{\max}$  of the active ingredient in the specific medium at pH 5.5 and 37°C. The quantity of active ingredient released is then determined as a function of time in the dissolution medium. At each time t, a sample of volume  $V_p$  is taken from the solution under study, then diluted with a medium of the same pH in a 10 cm<sup>3</sup> volumetric flask. The optical density (O.D.) of the diluted solution is then measured on the instrument calibrated at the corresponding wavelength (the dilution volume is chosen to obtain an O.D. value within the linear range of the calibration curve).

The following equation is used to calculate the mass of the liberated active agent (mt) at each instant t:

$$m_t = \text{DO} \cdot V_p \cdot V_f / \epsilon V_F l$$

$V_p$  = taken volume (1 mL).

$V_f$  = volume of solution after dilution (10 mL).

$V_F$  = volume of residual solution in the flacon following each precipitation.

$\epsilon$ : the specific absorption coefficient.

$l$ : length of quartz cell (1cm).

## II.9. BIOLOGICAL STUDY:

Medicinal plants exhibit significant biological activities such as antibacterial, antifungal, and cytotoxic properties [15,16]. These plants contain bioactive compounds that contribute to their medicinal properties and therapeutic applications [17]. Studies have shown that extracts from plants like *Acorus calamus*, *Artemisia annua*, and *Euphorbia helioscopia* demonstrate promising antibacterial and antifungal activities against various pathogens [18]. Additionally, research highlights the effectiveness of medicinal plant extracts from *Catharanthus roseus*, *Bixa orellana*, *Azadirachta indica*, and *Dracaena sanderiana* in combating bacterial and fungal infections [19].


The utilization of medicinal plants for their biological activities is rooted in their traditional use as natural remedies, offering a safer and cost-effective alternative to synthetic drugs. These findings underscore the importance of exploring the bioactive potential of medicinal plants for developing novel therapeutic interventions.

Antibacterial activity refers to the ability of a substance to either kill bacteria (bactericidal) or inhibit their growth (bacteriostatic) [20]. Various antimicrobial agents, including antibiotics, and plant extracts, exhibit antibacterial activity [21, 22]. For instance, plant extracts like *Imperata cylindrica* have demonstrated potent antibacterial activity against pathogens like *Staphylococcus aureus* and *E. coli*.

In our study, we examined the antibacterial activity of creams prepared and our aqua extracts in vitro against various bacterial strains, including *Staphylococcus aureus*, *Escherichia coli*, *Candida albicans* and *Pseudomonas aeruginosa*. To ensure their purity, these strains were seeded on selective media. The strains were then grown in Mueller Hinton (MH) agar medium and incubated at 37°C for 24 hours to obtain a young culture (24 hours old). Strain identification was carried out by **the laboratories of the Faculty Material Sciences, Ibn Khaldoun University.**

**Table II.10:** The tested bacterial stains.

| Reference strains | <i>Escherichia coli</i> | <i>Pseudomonas aeruginosa</i> | <i>Staphylococcus aureus</i> | <i>Candida albicans</i> |
|-------------------|-------------------------|-------------------------------|------------------------------|-------------------------|
| Code              | ATCC 25922              | ATCC 27853                    | ATCC 25923                   | ATCC 10231              |
| Gram              | Negative (-)            | Negative (-)                  | Positive (+)                 | Positive (+)            |



**Figure II.22:** Bacterial strains.



**Figure II.23:** Mueller Hinton Medium.


20 milliliters per Petri dish were aseptically prepared for the super cooled Mueller Hinton (MH) agar culture media. Let the medium cool and become solid on the surface of the work. After soaking a sterile swab in the bacterial solution, twist and press the swab firmly against the tube's inner wall to squeeze out any extra liquid. Using the swab, create tight ridges from top to bottom over the whole agar surface. Do this twice more, rotating the Petri dish through 60 degrees each time to cross the ridges and remembering to move the swab around on itself. Complete the seeding process by swabbing the agar's perimeter. As many Petri plates were inoculated, refill the swab was always necessary.

The analysis was released for the extracted plants and the prepared whites and their fresh creams.

For the extracted plants, discs are prepared by compression and were dropped on the Muller-Hinton surface while for the creams and their whites, they are placed in the created holes in Petri dishes. Each test was release in triplicate.



**Figure II.24:** Preparation of the petri dishes.



**Figure II.25:** Methods of product's statement.

Petri dishes were incubated at 37°C for 18 to 24 hours. Then the results were reading by measuring the diameter of the transparent zone of inhibition against the test microorganisms using a caliper.



**Figure II.26:** Bacterial dishes after incubation.

## **II.10. Minimal Concentration of Inhibitory**

The minimum inhibitory concentration (MIC) is defined as the lowest concentration of an antimicrobial agent capable of effectively inhibiting bacterial growth [23]. Research has shown that various natural extracts exhibit antibacterial properties at specific concentrations, making them potential alternatives for combating bacterial diseases in aquaculture [24, 25]. Determining how well these extracts prevent bacterial growth requires knowing their minimum inhibitory concentration.

### **II.10.1. Principe of the method:**

The minimum inhibitory concentrations (MICs) of plant extracts were initially determined using Mueller–Hinton broth microdilution . MIC determination was performed by a serial dilution technique using 96-well microtiter plates. Plant extract (100  $\mu$ L) was placed into the well/plate. Then, 100  $\mu$ L bacterial cell suspensions were placed in each well/plate. Microplates were incubated for 24 h at 37 °C. The lowest concentrations without visible growth completely inhibited the bacteria (MICs) [26].






**Figure II.27:** Microplates for the MCI method.

### II.10.2. Protocol of the MCI Determination:


The extracted solutions were prepared to be tested with a mass of 0.01g of extract solubilized in 1mL of DMSO. The spaces of the microplates were filled with 100 $\mu$ L of Mueller-Hinton microdilution broth (96 spaces). The prepared solution was deposited in the first space and homogenized with micropipettes, then a second dilution was released from the first solution. The process was repeated as well as to fill the 10 spaces. 20 $\mu$ Ls were deposited in all spaces and the last space was considered as reference.



(1) : 100 $\mu$ L of Mueller-Hinton microdilution broth



(2) : Extract plants (100  $\mu$ L) were placed into the microplate.



(3): Addition of 20  $\mu$ L of bacterium  
medium



(4) : Microplates were incubated for 24 h  
at 37 °C hours


**Figure II.28:** Protocol of the MCI determination.

## II.11. Antioxidant activity

### II.11.1. Principle of the method:

Numerous methods are currently used to assess this activity; the antioxidant activity of a compound corresponds to its ability to resist oxidation [27]. The 2,2-diphenyl-1-picrylhydrazyl radical has been widely used to study the free radical scavenging activity of various extracted plants. The chemical compound 2,2-diphenyl-1-picrylhydrazyl (DPPH) was one of the first free radicals used to study the structure-antioxidant activity of phenolic compounds [28]. It possesses an unpaired electron on one atom of the bridge, the reduction of this radical is accompanied by its from the characteristic violet color of the DPPH solution to the yellow color measurable spectrophotometrically at 514-518 nm [29].

DPPH is a stable free radical with a dark purple dark violet color that turns pale yellow when reduced in the presence of an antioxidant [30].



**Figure II.29:** Structure of the DPPH radical and its reduced form.

### II.11.2. Antioxidant experiment:

The antioxidant activity of the extracted plants and the standard was assessed on the basis of the radical scavenging effect of the stable 1, 1-diphenyl-2-picrylhydrazyl (DPPH)-free radical activity by modified method. The diluted solutions of the extracted plants were prepared in methanol 0.002% of DPPH which was prepared in methanol. 1.5 mL of this solution was mixed with 1 mL of prepared and standard solutions separately. These mixtures were kept in dark for 30 min and optical density was measured at 517 nm. Methanol (1 mL) with DPPH solution (0.002%, 1.5 ml) was used as blank, and the same method of ascorbic acid at different concentration was used as reference [31].


The difference in the absorbance between the test solution and the control (DPPH in methanol) was calculated and expressed as scavenging percent or inhibition percent of DPPH radical. The inhibition percent of DPPH radical was calculated by using the equation; Percent inhibition of DPPH.

$$I = \left( \frac{A_c - A_s}{A_c} \right) * 100$$

Whereas: I %: Trapped DPPH rate or inhibition rate.

As: Is the absorbance of the test solute

Ac: Is the absorbance of the control.



**Figure II.30:** Antioxidant activity protocol.

### II.11.3. IC<sub>50</sub> estimation:

IC<sub>50</sub> is defined as the concentration of the sample required to reduce 50% of the DPPH radical, at IC<sub>50</sub>s are determined graphically by linear regression of the plotted graphs (inhibition rate as a function of different concentrations).

### II.12. Anti-inflammatory activity

The anti-inflammatory activities of the crude and fractionated plant extracts were determined using a modified version of the BSA assay reported by Williams et al.<sup>11</sup> BSA solution (0.4%, w/v) was prepared in Tris Buffered Saline, pH 6.4 at 25 °C. Stock solutions of each plant extract were prepared at a concentration of 50 µg/mL or 0.005%, w/v. Respective aliquots of 5.0 µL, 10 µL and 20 µL representing concentrations of 200 µg/mL and 1.00 µg/mL of the stock solutions were added to test tubes containing 1 mL of 0.4%, w/v BSA buffer solution. Both negative (methanol) and positive (diclofenac) controls were assayed in a similar manner. The solutions were then heated in a water bath at 72 °C for 10 minutes, and cooled for 20 minutes under laboratory conditions. The turbidity of the solutions (level of protein precipitation) was measured at 660 nm in a Hatch Spectrophotometer using an air blank. The experiments were conducted in duplicate and the mean absorbance values were recorded. The percentage inhibition of precipitation (protein denaturation) was determined on a percentage basis, relative to the negative control using the following equation: [32].

- Anti-Denaturation Activity% =  $\frac{(\text{Absorbance of control} - \text{Absorbance of sample})}{(\text{Absorbance of control})} \times 100$ .

- Anti-Denaturation Activity% = % Inhibition of Protein Denaturation = % Anti-inflammatory Activity.

### II.13. In vivo tests

In this part, two types of experience are described. The first is to test the anti-inflammatory activity whoever the second is to test the ability to progressive the hair of rats.

- **First experience:**

The first in vivo experiment was conducted on a total of twenty-four (24) Wistar rats, sourced from the Pasteur Institute of Algiers. The rats, aged 4 months with weights ranging from 220 to 322g (mean weight=261±5g) at the beginning of the study, had undergone a 4-month adaptation period at the Hygiene and Animal Pathology Laboratory, Tiaret. Where they were housed at room temperature and provided with ad libitum access to food and water.

Rats are housed individually in plastic cages with wood shavings bedding, which is changed daily. Each cage was fitted with a tag bearing the name of the batch corresponding to the treatment. The rats were divided into 8 distinct groups each group consist 3 males.



**Figure II.31:** Presentation of rat's group.

After that the rats were fasted for 18 hours, measurements of the right paw were taken then, an injection of 1 mL of a 1% formaldehyde solution is administered into the plantar aspect of the right paw to induce edema.





**Figure II.32:** Injection to induce edema.

We premeasured the paw's diameter two hours after the injection since we saw redness and swelling at the injection site.



**Figure II.33:** Apparition of edema and their measurement.

The edematous paw was treated by prepared galenic forms. The treatments were given to test rats for 5 days with free access to water and food each day, the description of the groups is summarized in the next table.

**Table II.11:** Description of rat's group of the first experience.

| Number of groups      | 1           | 2            | 3           | 4            | 5            | 6            | 7            | 8                |
|-----------------------|-------------|--------------|-------------|--------------|--------------|--------------|--------------|------------------|
| Number of Rats /group | R16, R2, R8 | R1, R14, R19 | R5, R7, R22 | R4, R11, R21 | R3, R12, R20 | R8, R15, R24 | R9, R10, R23 | R13, R6, R17     |
| Treatment             | L1          | L2           | L3          | L4           | L5           | L6           | L7           | Positive control |


Every day prior to the tendering of prepared galenic forms, the diameter of the rats' paws is measured with a digital caliper to track the progression of the edema.

- ***Second experience:***

The second in vivo experiment was conducted on a total of ten (10) males Wistar rats, With the same conditions, Rats were housed individually in plastic cages with wood shavings bedding, which is changed daily. Each cage was fitted with a tag bearing the name of the batch corresponding to the treatment. The rats were divided into 5 distinct groups each group consist 2 males.

After placing the mice in their designated places, we shaved the back of the mouse with an area of approximately 9 square centimeters.





**Figure II.34:** Hair removal zone.

**Table II.12:** Description of rat's groups for the second experience

| Number of groups      | 1                | 2           | 3                         | 4           | 5               |
|-----------------------|------------------|-------------|---------------------------|-------------|-----------------|
| Number of Rats /group | R1, R2           | R1, R2      | R1, R2                    | R1, R2      | R1, R2          |
| Treatment             | Positive control | Basic Cream | Basic Cream + Thy shampoo | Thy shampoo | Thy Cam shampoo |

## *Experimental references*

1. Azza, Z., & Oudghiri, M. (2015). In vivo anti-inflammatory and antiarthritic activities of aqueous extracts from *Thymelaea hirsuta*. *Pharmacognosy research*, 7(2), 213-218.
2. Ankita Jain, DV Rao, Amla Batra and Amita Jain, *Elixir Appl. Botany*, 2014, 77, 28867-28871.
3. ROOPALATHA C., VIJAY M., *International Journal of Pharmacy and Pharmaceutical Sciences*, 2013, 4(5), 629-634.
4. Prakash C.V.S, *Int. J. Res. Chem. Environ*, 2014, 1(1), 1-18.
5. K. Sahira Banu, Dr. L. Cathrine, *International Journal of Advanced Research in Chemical Science*, 2015, 2(4), 25-32.
6. Jeevarathinam A., Muthulakshmi P., *Ijppr.Human*, 2017, 8 (3), 232-241.
7. Hajoori M, Naik M, Naik K, Desai S, *International journal of pharmacological screening methods*, 2014, 4, 26-41.
8. Ahmed S. A., Abood N. H., Al-Janabi A. A., *Tech. J.*, 2013, 31(3), 1-5.
9. Nirmala Phuyal, Pramod Kumar Jha, Pankaj Prasad Raturi, and Sangeeta Rajbhandary, Total Phenolic, Flavonoid Contents, and Antioxidant Activities of Fruit, Seed, and Bark Extracts of *Zanthoxylum armatum* DC. *Hindawi the Scientific World Journal*, Vol2020, pp. 1-7, 2020.
10. A. Marsoul, M. Ijjaali, F. Elhajjaji, M. Taleb, A. Boukir, *Mater. Today Proc.* 27 (2020) 3193–3198.
11. D. Kim, O. Chun, Y. Kim, H. Moon, C. Lee, *J. Agric. Food Chem.* 51 (2003) 6509d
12. CB Singh, MC Devi, DS Thokchom, Sengupta M, AK Singh, Phytochemical screening, estimation of total phenols, total flavonoids and determination of antioxidant activity in the methanol extract of *Dendrobium denudans* D. Don stems; *Journal of Pharmacognosy and Phytochemistry* 2015;4(4):06-11.
13. Sun.B, Richardo-da-Silvia JM, Spranger I. Critical factors of vanillin assay for catechins and proanthocyanidins [J]. *J Agric Food Chem*, 1998, 46: 4267-4274.


## *Experimental references*

- 14.** Madhav Poudel and Meena Rajbhandari, Phytochemical Analysis of *Ampelopteris Prolifera* (Retzius) Copeland; Nepal Journal of Science and Technology; (NJST) (2020), 19(1): 78-88.
- 15.** Francisco, Les., Guillermo, Cásedas., Víctor, López. (2021). Bioactivity of Medicinal Plants and Extracts. *Biology*, doi: 10.3390/BIOLOGY10070634.
- 16.** Abdulaziz, M., Al-Othman., Iqbal, Hussain., Hamayun, Khan., Muneeb, Ur, Rehman., Ahmed, A., Abdeltawab., Riaz, Ullah., Rohullah., Shumaila, Noor., Muhammad, Talha. (2012). Phytochemical analysis and biological activities of selected medicinal plants. *Journal of Medicinal Plants Research*.
- 17.** Abbas, Zare, Banadkoki., Ebrahim, Kouhsari., Nour, Amirmozafari., Maryam, Roudbary. Mohammad, Reza, Boujary, Nasrabadi. (2018). Antibacterial, antifungal and cytotoxic activities of some medicinal plants against multidrug resistance pathogens. *Reviews in Medical Microbiology*.
- 18.** Ushie O. A, et.al., Bioactivity of Active Extracts of Leaf of *Bauhinia Racemosa* *International Journal of Clinical Chemistry and Laboratory Medicine* 2019, 5(2): 19-24.
- 19.** Shikha, Sharma., Neha, Dangi., P., K., Goyal., Jeyabalan, Govindasamy. (2023). The activity of several medicinal herbs such as anti-inflammatories, anti-pyretic, analgesics and antioxidants. *Current Indian Science*.
- 20.** Marius, Belmondo, Tincho., Thureyah, Morris., Mervin, Meyer., Ashley, Pretorius. (2020). Antibacterial Activity of Rationally Designed Antimicrobial Peptides. *International Journal of Microbiology*.
- 21.** V., Parkavi., M., Vignesh., K., Selvakumar., Mohamed, J., Muthu., J., Joysa, Ruby. (2012). Antibacterial activity of aerial parts of *imperata cylindrica* (l) beauv. *International Journal of Pharmaceutical Sciences and Drug Research*.
- 22.** Kehinde, Tope, Adegbehingbe., Soji, Fakoya., Marcus, Oluyemi, Bello., Charles, Ayodeji, Osunla., Samson, Olumide, Akeredolu. (2019). Antibacterial Activity of Bacteria Isolated from Fermenting Cocoa Water against Some Pathogenic Bacteria. *International Journal of Current Microbiology and Applied Sciences*.

## *Experimental references*

- 23.** Rusmawanto, Rusmawanto., Arief, Prajitno., Ating, Yuniarti. (2019). Minimum Inhibitory Concentration of Marine Microalgae *Dunaliella salina* on Fish Pathogenic Bacteria *Edwardsiella tarda*. doi: 10.21776/UB.RJLS.2019.006.02.1.
- 24.** Grace, Ramena., Yathish, Ramena., William, Duke., Nitin, Challa. (2018). Determination of Minimum Inhibitory Concentrations of Antimicrobial Herb and Spice Extracts against *Flavobacterium columnare*.
- 25.** Robert, F., Moran., Emery, N., Brown. (1997). The Smallest Concentration. *Clinical Chemistry*, doi: 10.1093/CLINCHEM/43.5.856.
- 26.** Wikaningtyas, P., & Sukandar, E. Y. (2016). The antibacterial activity of selected plants towards resistant bacteria isolated from clinical specimens. *Asian Pacific Journal of Tropical Biomedicine*, 6(1), 16-19.
- 27.** Rice-Evans CA, Miller NJ, Bolwell PG, et al. (1995) The relative antioxidant activities of plant derived polyphenolic flavonoids. *Free Radical Res* 22: 375–83.
- 28.** Brand-Williams W, Cuvelier ME, Berset C (1995) Use of a free radical method to evaluate antioxidant activity. *Lebensmittel-Wissenschaft und Technologie* 28 : 25–30.
- 29.** Bentabet, N., Boucherit-Otmani, Z., & Boucherit, K. (2014). Composition chimique et activité antioxydante d'extraits organiques des racines de *Fredolia aretioides* de la région de Béchar en Algérie. *Phytothérapie*, 12(6), 364-371.
- 30.** NABTI, B., SISALAH, I., TAIBAOU, K., & BOURKAIB, S. (2023). Evaluation of antioxidant and antimicrobial activities of polyphenols from leaf extracts of a spontaneous *Phlomis* from Northern Algeria. *Batna J Med Sci*, 10(1), 96-101.
- 31.** Khalaf, N. A., Shakya, A. K., Al-Othman, A., El-Agbar, Z., & Farah, H. (2008). Antioxidant activity of some common plants. *Turkish Journal of Biology*, 32(1), 51-55.
- 32.** Williams L., Vasquez E., Milan P., Zebitz C., Kraus W. In vitro anti-inflammatory and anti-microbial activities of phenylpropanoids from *Piper betle* (Piperaceae). In proceeding of the Phytochemical Society of Europe: Natural products in the new millennium: Prospects and industrial application. AP Rauter, PB Palma, J. Justino, Araujo ME, Santos SP(Eds). Kluwer Academic Publisher, Dordrecht. The Netherlands.2002;74:221-27



The image features a blue-tinted background with a grid pattern. Overlaid on this are various handwritten notes and diagrams in a lighter blue color. A large magnifying glass with a black frame is positioned in the center-right, focusing on a white oval area. Inside this oval, the text 'Chapter III Results and discussions' is written in a bold, italicized black font. The magnifying glass's handle is visible at the bottom right corner.

***Chapter III  
Results and  
discussions***



### III.1. INTRODUCTION:

This section will cover the results and the characterization of prepared galenic forms and plant extracts. Bacterial tests and medication release investigations will also be covered. In vivo studies were used to confirm their activity.

### III.2. CHARACTERISATION OF EXTRACTS

#### III.2.1 PHYSICO-CHEMICAL CHARACTERIZATION OF EXTRACTS PRODUCT:

- **Yield:** the yield of our experiment is calculated compared with the initial quantity of the plant as described the next equation.

$$\text{Yield} = (\text{the mass of obtained extract} / \text{plant's initial mass}) * 100$$

The calculated values were summarized in the table III.1. The obtained values were compared with previous research and they showed interesting differences in values when organic solvents were used as extracted medium [1].

**Table III.1:** The calculated yield.

| Plants | yield % |
|--------|---------|
| Thy    | 15.73   |
| Cam    | 14.254  |
| Cur    | 0.03    |

The yield of the extracted curcumin was little and the obtained mass was analyzed in FTIR and the solubility test.




Figure III.1: Extraction yields.


- Solubility:** it's important to test the solubility of our extracts product to choose initially the composition of our galenic forms. The obtained results are summarized in the table III.2.

Table III.2: The solubility's test.

| Product          | Thy   | Cam   | Cur   |
|------------------|-------|-------|-------|
| Water            | +++++ | +++++ | ----- |
| alcohol          | ----- | ++++  | ----- |
| propylene glycol | ----- | ----- | ----- |
| paraffin oil     | ----- | ----- | +++++ |
| Vaseline oil     | ----- | ----- | +++++ |
| coco betaine     | ++++  | +++++ | ----- |
| DMSO             | +++   | +++   | +++++ |



- **Refractive index:** the refractive index was determined using the refractometer and the obtained values are 1.338 for the chamomile extract and 1.337 for the thymeleae hirsute extract.
- **pH:** the pH of the extracted solution of the studied plants are measured and the obtained values are 5.5 and 5 of thymeleae and chamomile respectively. The pH of extracted curcumin in acetone is 4.5.
- **FTIR analysis:** FTIR gives an idea about the composition of the analyzed product. The characterized bands were compared to another work [2]. The obtained IR spectra confirm the presence of characteristic bands corresponding to determined compounds in phytochemical analysis. The spectra are shown in the figures below and the bands values are reported in the table III.3:



|   |                    |                                    |            |
|---|--------------------|------------------------------------|------------|
| C:\OPUS_7.2.139.1294\MEAS\Sample description.4833 | Sample description | Instrument type and / or accessory | 18/02/2024 |
|---|--------------------|------------------------------------|------------|

Seite 1 von 1

Figure III.2: Infrared spectrum of thymeleae.




Figure III.3: Infrared spectrum of chamomile.




Figure III.4: Infrared spectrum of curcumin.

**Table III.3:** Characteristic bands of extract plants.

| Plants           | Bands  |
|------------------|--|
| <b>Thymelaea</b> | 1030.41 cm <sup>-1</sup> correspond to C-O function.<br>1598.21 cm <sup>-1</sup> correspond to C=C function.<br>2846.77 cm <sup>-1</sup> correspond to unsaturated C-H function of aldehyde.<br>2914.77 cm <sup>-1</sup> correspond to saturated C-H function.<br>3263.33 cm <sup>-1</sup> correspond to O-H function. |
| <b>Chamomile</b> | 1030.01 cm <sup>-1</sup> correspond to C-O function.<br>1402.27 cm <sup>-1</sup> correspond to OH function.<br>1585.53 cm <sup>-1</sup> correspond to C=C aromatic function.<br>2921.31 cm <sup>-1</sup> correspond to saturated C-H function.<br>3250.20 cm <sup>-1</sup> correspond to O-H function.                 |
| <b>Curcumin</b>  | 1268.29 cm <sup>-1</sup> correspond to C-O function.<br>1577.72 cm <sup>-1</sup> correspond to C=C aromatic function.<br>2920.28 cm <sup>-1</sup> correspond to C-H function.<br>3338.90 cm <sup>-1</sup> correspond to O-H function.  |

The first two spectra present bands around 3263–3250 cm<sup>-1</sup> in a row corresponding to the OH stretching vibrations characteristic of organic acids. Additionally, there are two medium-intensity bands at 2914 cm<sup>-1</sup> and 2921 cm<sup>-1</sup>, indicative of the symmetric C-H stretching vibrations of aliphatic CH<sub>2</sub> and CH<sub>3</sub> groups, respectively. A band in the range of 1598 to 1585 cm<sup>-1</sup> are attributed to the C=C stretching vibrations of aromatic rings and the bands in the vicinity of 1413-1402 cm<sup>-1</sup> are assigned to OH deformation vibrations. A strong band at approximately 1030 cm<sup>-1</sup> is attributed to phenolic C–O stretching, however the bands in the range of 616 cm<sup>-1</sup> are associated with the deformation vibrations of unsaturated CH bands. The presence of these characteristic absorption bands confirms the richness of the leaf extracts in phenolic compounds in the two studied extracts.

The FTIR spectrum of Cur shows a weak band at  $3338.80\text{ cm}^{-1}$ , attributed to the O-H stretching. The band appearing at  $1626.11\text{ cm}^{-1}$  corresponds to the stretching vibration of the aromatic C=C band and a moderately intense band at  $1577.72\text{ cm}^{-1}$  is associated with benzene ring stretching. The sharp and intense band at  $1508.69\text{ cm}^{-1}$  corresponds to the C=O and C=C vibrations of Cur. The band located at  $1268.29\text{ cm}^{-1}$  is attributed to the C-O bond vibration.

### III.2.2. PHYTOCHEMICAL ANALYSIS:

The phytochemical evaluation of various phytoconstituents in the studied extracts were graded as high (+++) and nil (-) based on the visual colored reaction product of the test.

**Table III.4:** Result of phytochemical tests.


| test        | aqua extract of Thy | aqua extract of Cam |
|-------------|---------------------|---------------------|
| tannin      | +++                 | +++                 |
| saponins    | +++                 | +++                 |
| quinones    | +++                 | +++                 |
| poly phenol | +++                 | +++                 |
| flavonoids  | +++                 | +++                 |
| steroids    | -                   | -                   |
| alkaloids   | +++                 | +++                 |
| coumarins   | +++                 | +++                 |

The employed phytochemical tests indicate the presence of all tested products except the steroid product. For this the obtained products were analyzed and the result will be presented.

### III.2.3. QUANTITATIVE ANALYSIS

#### ✚ Determination of total polyphenols

The total polyphenol content of the different extracts is determined using the regression equation  $y=0.001x-0.031$  from the calibration curve of gallic acid which is classed in annex. The results are expressed in milligrams of gallic acid equivalents per gram of dry matter (mg GAE/g) [3].




**Figure III.5:** Total polyphenol content of extracts.

Thy and Cam had a total phenolic content of 10.61 mg GAE/mg extract and 12.48 mg GAE/mg respectively, these values are lower than obtained values with alcoholic solvent in previous works [4].

#### ✚ Determination of total flavonoids

The total flavonoids content of the different extracts is determined using the regression equation  $y=0.271x+0.0004$  from the calibration curve of quercetin. The results are expressed in milligrams of quercetin equivalents per gram of dry matter (mg GAE/g) [5].




**Figure III.6:** Flavonoid content of various extracts.

The histogram showed the total content flavonoids in thy and cam extracts. Thy extract had a total flavonoid of 18.11 mg QE/mg and the cam extract had 15.85 mg QE/mg. The obtained values are compared to literature, and they are close [6].

#### **✚ Determination of total tannin content**

The tannin content is calculated using the calibration curve of catechin,  $y = 0.00076 x - 0.0076$  [7]. It is expressed in milligrams of catechin equivalents per grams of extract (mg CE/g).

















**Figure III.7:** Tannin content of different extracts.

The studied extracts showed the tannins content of 38.08mg/g for cam and 46.71 mg/g for thy. The use of water as extract solution gives a notable value of chemical compounds as previously determined.

### III.3. CHARACTERISATION OF GALINICS FORMS:


- **Color:** our creams adapted the color of the extract.
- **Oder:** the odor of our creams is fragrant from the paraffin oil and the other corposants.
- **consistency:** All ointments display a semi-solid texture.
- **Stability:** Ointments maintain stability at 27°C and liquefy at 35°C.
- **Homogeneity:** Our ointments demonstrate excellent uniformity upon spreading several samples on a flat surface.
- **Centrifugation:** we do the test of separation with the centrifugation method. So, the centrifugation was release at 5 min and 30 min, the results are shown in the table III.5:

**Table III.5:** Result of centrifugation of creams.

| Creams | L1  | L2  | L3  | L4  | L5   | L6  | L7  |
|--------|---|---|---|---|--|---|---|
| Time   |   |   |   |   |  |   |   |
| 5 min  |  |  |  |  |  |  |  |
| 30 min |  |  |  |  |  |  |  |



- **Spreading:** the creams were passed over a glass to identify their consistence and their morphology when using on the skin. The represented photos are summarized in figure III.8.



**Figure III.8:** Spreading of some creams. (A: Cur cream, B: Cam cream and C: Thymelaea cream).

- **pH Level:** Consistency is observed in the pH levels across all formulation so that we obtain the following results:  
The pH of prepared galenic forms was measured and the obtained values were classified in the table III.6., they present a pH close to the skin pH.

**Table III.6:** The galenic forms pH.


| Cream | pH  |
|-------|-----|
| L1    | 5.5 |
| L2    | 5.5 |
| L3    | 5   |
| L4    | 5   |
| L5    | 5.5 |
| L6    | 5.5 |
| L7    | 5   |

### III.4. RELEASE STUDY


The release of extracted plants from various prepared formulations has been evaluated by measuring the amount of PA that is released over time using a UV-visible spectrophotometer. The results are presented in the figures III.9-III.15.

The obtained data indicates a low release percentage, not exceeding 50% for the majority of prepared galenic forms. This is primarily justified by the lipidic composition of the prepared galenic forms, which makes the liquid's penetration (pH=5.5) into the interior of the galenic form difficult. However, the lipid-rich environment of the skin facilitates their penetration.


The obtained release curves indicate that the release rate is highest at the beginning and then gradually decreases that is explained by the slow diffusion of the active ingredient. After 50mn, a continuous release pallier achieved the diffusion.




**Figure III.9:** Liberation curve of Cur from L1.




**Figure III.10:** Liberation curve of Cur from L2.



**Figure III.11:** Liberation curve of Cam from L3.



**Figure III.12:** Liberation curve of Cam from L4.



**Figure III.13:** Liberation curve of active agents from L5.




Figure III.14: Liberation curve of active agents from L6.





Figure III.15: Liberation curve of Thy from L7.

### III.5. BACTERIAL TEST


The activity of the studied extracts and their formulations were assessed based on the presence or absence of bacterial growth inhibition. The presence of antibacterial activity was indicated by clear zones around the discs soaked with the extracts, while the absence of inhibition was reflected by the absence of such halos around the wells. This evaluation was conducted after 24 hours of incubation at 37°C. The obtained results are presented in the following figures:



**Figure III. 16:** Results of antibacterial activity against different strains (a, b, c, d: extracted plants, e: L5 and L6, f: L3 and L4).



The antibacterial activity tests revealed different activities against the various tested strains. The inhibition zones and their diameters are presented in the tables (Annex II.4) and in the figures below.



**Figure III.17:** Inhibition diameters of L1.0, L2.0, L1, L2, and the cur extract against the studied strains.

According to the figure (III.17), it is evident that cur extract demonstrates antifungal activity; however, when formulated as L1, it exhibits an antibacterial activity against *E. Coli* and *Staph*,


This can be explained by the associative effect between L1.0 and cur extract against these strains, but for the *Candidat Albicans*, the antagonist effect is noted.

The L2 ointment has shown a very good antifungal and bacterial activity of the core extract compared to L1, the composition of the L2 ointment has a considerable antifungal effect and an important antibacterial effect once associated with cur extract.


According to the figure III.18, the cam extract presents an important activity against all studied strains. For L3, the antibacterial activity was preserved after formulations, but when comparing the ointment L3 and L3.0, an associative effect is noticed for the *pseudo* and an antagonist effect for the *E. Coli* and *Candidat*.

On the other hand, For the L4 and L4.0 ointment, an associative effect is noted for the *Pseudo*, and an antagonistic effect for the *E. coli*, *staph* and *Candidat strains*.

The formulation of cam (L3, L4 ointments) gave an antifungal activity, which is not found in the base ointment L3.0 and L4.0.



**Figure III.18:** Inhibition diameters of L3.0, L4.0, L3, L4, and the cam extract against the studied strains.




**Figure III.19:** Inhibition diameters of L5.0, L6.0, L5, L6, and the cam, cur extracts against the studied strains.

By comparing the biological activity of curcumin and the cam extracts with the ointment L5 and L6 loaded in different percentages, we note, that the two ointments have a good biological activity once the extracts are combined.

Comparing the composition of L5 and L6 formulation, the antifungal activity is accentuated when cur present 2% (L5) and the bacterial activity is important against *pseudo* and *staph* when cam present 2% (L6).

It is also noted that the base ointment L6.0 has a very good biological activity without being associated with any active extract.



**Figure III.20:** Inhibition diameters of L7.0, L7, and the thy extract against the studied strains.

The L7 ointment loaded with the thy extract has an average antibacterial activity in comparison with the base ointment which has an important antifungal activity. The combination of the base ointment with the extract canceled the antifungal activity and the bacterial activity against *staph* of the base ointment.

The thy extract alone has a very low biological activity which will give an associative effect when it's formulated to L7 ointment, vis-à-vis the *pseudomonas* strain and *E. Coli* strain.

### III.6. MINIMAL CONCENTRATION OF INHIBITORY

For the evaluation of the effectiveness of the prepared formulations, it is necessary to determine the minimum inhibitory concentration. The obtained results of well plates and the tested concentration are summarized in the next tables.





**Figure III.21:** MIC results for Thymelaea extract.



**Figure III.22:** MIC results for Chamomile extract.


Thy extract has the most important minimum inhibitory concentration noted for *staph* which is 12.5mg/mL and for the other strains the MIC is 6.25mg/mL, this can be explained from the resistance of the parity of the heterogeneous *staph* strain which consists of proteins that are polymers difficult to degrade by the extracts. By comparing the non-rigid parity of gram strains, these observations are similar for the extract of turmeric as far as chamomile is concerned, the

minimum inhibition of concentration is 3.12mg/mL for the entire strain. The least values of MIC of tested extracts correspond to the cam so, the most effective extract is chamomile, a lower MIC.


### III.7. ANTIOXYDANT ACTIVITY

The obtained curve of Cur demonstrates a negative value which show that Cur hasn't an antioxidant activity.

Thymeleae extract reduces the stable free radical 2,2-diphenyl-1-picrylhydrazyl (DPPH) to yellow-colored diphenylpicrylhydrazine with an IC<sub>50</sub> of 13.68 mg/mL showing an antioxidant activity, greater than ascorbic acid and Cam activities which represent 44.8 mg/mL and 74.24mg/mL respectively. It can be concluded that the most effective extract is that of thymeleae.



**Figure III.23:** Inhibition percentage of free radical DPPH in function of concentration of Cam extract.



**Figure III.24:** Inhibition percentage of free radical DPPH in function of concentration of Thy extract.



### III.8. ANTIINFLAMMATORY ACTIVITY

The maximum inhibition and denaturation effect of the THYML extract at the concentration of 100 $\mu$ g/ mL was 65.45%, whereas the diclofenac standard at 100  $\mu$ g/mL showed a 98.19% inhibition of the denaturation of the SAB. The correlation between the concentration and the percentage of inhibition showed that the inhibition capacity of the denaturation of the BSA that the extract of thy has a greater anti-inflammatory effect than that of chamomile which remains weaker than diclofenac.

**Table III.10:** % inhibition of denaturation of extracted plants.

| Concentration  | Extracted plants       | % Inhibition of denaturation |
|----------------|------------------------|------------------------------|
| 100 $\mu$ g/mL | Thy                    | 65.45%                       |
| 100 $\mu$ g/mL | Cam                    | 39.97%                       |
| 100 $\mu$ g/mL | Diclofenac (reference) | 98.19%                       |

### III.9. IN VIVO TEST

#### First experience:

In order to assess the anti-inflammatory properties of prepared galenic forms, it was tested on Wistar rats that had been experimentally injected with 1% formol at the level of their right paw.




**Figure III.25:** paw of rat before treatment.



**Figure III.26:** rat paw after treatment.

The evolution of the patten diameters is illustrated in figure III.26.



**Figure III.27:** Evolution of the patten diameters.

The oedema appears within the first hour of the first hour of the 1% formol injection.

The observations indicate that batch 8 (rats injected with 1% formol without treatment) presents a low decrease of the paw's diameter due to the immune system of the tested rats without returning to normal state at the end of the experiment.

The rats treated with prepared galenic forms showed a significant reduction in oedema and were back to its normal state at the sixth day. On the other hand, rats given L2 and L3, which contains Cur1%+ZnO 1% and Cam 1% respectively, their paws returned to normal state at the fifth day and the increasing is notable at the second and third days.

The galenic forms containing Cam 1% (L3 and L5) show an important reducing in oedema compared to the others with 1,5% and 2% (L4 and L6).

The L7 presents a low decreasing in paw's diameter compared to the other formulations, this is justified by the power of anti-inflammatory activity of Cam, Cur compared to the Thy extracted plants.

### Second experience:






In order to evaluate the activity of thymeleae extract on the growth of hair, basic cream and two shampoo were prepared and tested on different batches.

As demonstrated in summarized photos, the growth of hair is important in treated batches compared to the control batch.





















In the fourth week, the hair's evolution of control batch achieves 15% and 35% in the fifth week. this incomplete growth is clear at the last photo.

In the fourth week, the use of basic cream with shampoo gives an important growth of hair accomplish to 80%, however the separately use of shampoo and basic cream gives 60% and 70% of evolution growth respectively. A thick hair for the treated batches was noticed at the end of experience.

**Table III.11:** The growth of hair over the 5 weeks for the control Batch.

| Weeks   | First week  | Second week   | Third week   | Forth week  | Fifth week  |
|---------|---|---|--|---|---|
| Batch   |  |  |  |  |  |
| Batch 1 |   |   |  |   |   |

**Table III.12:** The growth for hair over the 5 weeks for the treatment Batches.

| Weeks<br>Batches | First week  | Second week  | Third week  | Forth week  | Fifth week  |
|------------------|---|--|---|---|---|
| Batch 2          |    |    |    |    |    |
| Batch 3          |    |    |    |    |    |
| Batch 4          |   |   |   |   |   |
| Batch 5          |  |  |  |  |  |

## *Result and Discussion references*

1. ROOPALATA and VIJAY MALA 2013 International journal of pharmacy and pharmaceutical sciences
2. Morozhenko V., Infrared Radiation in Tech, 2012, 3 rd edition, Lavoisier, doi :10.5772/2031.
3. Ali-Rachedi, F., Meraghni, S., Touaibia, N., & Mesbah, S. (2018). Analyse quantitative des composés phénoliques d' ; une endémique algérienne Scabiosa Atropurpurea sub. Maritima L. Bulletin de la société royale des sciences de liège
4. (Mounira Merghem, Saliha Dahamna, Seddik Khennouf, Journal of Drug Delivery & Therapeutics. 2020; 10(4):108-111) (Khennouf Seddik, Benchiekh Dalila, Djidel Saliha, Dahamna Saliha, Amira Smain, Charef Noureddine, Baghiani Abderahmane, Harzallah Daoud, Arrar Lekhmici, Pharmacognosy Communications, 2013, 3(2), 35-40)
5. Bohui, P. S. G., Adima, A. A., Niamké, F. B., & N'Guessan, J. D. (2018). Etude comparative de trois méthodes d'extraction des flavonoïdes totaux à partir des feuilles de plantes médicinales : Azadirachta indica et Psidium guajava. Journal de la Société Ouest- Africaine de Chimie, 46, 50-58.
6. (Roumaissa Ounis, Fatima Benchikh, Smain Amira, Hassiba Benabdallah, Hind Amira, Walid Mamache, Bensouissi Chawki, Khaoula Hellal, Turkish Journal of Agriculture - Food Science and Technology, 2023, 11(6): 1161-1167). (G. Haghi, A. Hatami, A. Safaei, M. Mehran, Research in Pharmaceutical Sciences, 2014; 9(1): 31-37)
7. vinson JA, Dabbagh YA, SerryMM, al Plant flavonoids, especially tea flavonoide, are powerful antioxydants using an in vitro oxydation model for heart disease[j]Agric FoodChem,1995,43 :2800-2802



# *General conclusion*

## *General conclusion*

Medicinal plants have many established therapeutic uses because they can contain numerous active chemicals. For this purpose, this work concentrates on some extracted plants and their formulation into galenic forms (creams and shampoo).

Initially, the extracted of essential oil was proposed and since it was difficult to obtain, one moved on to plant extracts. These last were attained through hydro-maceration and they were characterized by UV, IR-spectrum, and their phytochemicals analysis were also studied and quantified. The results show that extracted plants using water gives a notable value of active chemical compounds.

Many galenic forms were prepared and they were tested to confirm their stability, the effectiveness of these creams and were analyzed by anti-bacterial, antioxidant, anti-inflammatory tests. The obtained fallouts were compared of extracted plants' results.

The extracted plants and their formulation present and important bacterial and antifungal activities with noticed difference between a formulation to other, this remark is justified by the antagonist and additional effects of some excipient or combination between extracted plants. Consuming the same used strains (*staph*, *pseudo*, *E. Coli* and *candida*), the minimal inhibitory concentration was determined, and the Thy extract has an MIC noted for *staph* 12.5mg/mL and for the other strains 6.25mg/mL. The least value of MIC is 3.12mg/mL correspond to the cam which present the most effective extract.

Thy extract present an antioxidant activity better than that cam extract, this is being noticed by the calculated IC<sub>50</sub> of 13.68 mg/mL and 74.24mg/mL of Thy and cam extracts respectively.

The anti-inflammatory in vitro shows that Thy extract has a greater anti-inflammatory effect than that of chamomile which remains weaker than diclofenac and the in vivo test illustrates that all creams present an anti-inflammatory activity demonstrating a difference of reducing oedema.



## *General conclusion*

Testing the prepared shampoo, the growth of hair is noted in treated batches compared to the control batch. the use of basic cream with shampoo gives an important growth of hair accomplish to 80%, however the separately use of shampoo and basic cream gives 60% and 70% of evolution growth respectively. A thick hair for the treated batches was observed at the end of experience.

As conclusion, the galenic forms (creams and shampoo) give important upshots in studied biological activities as so as their extract plants.




Figure 1: Spectrum of absorption of the aqua extract of Chamomile.




Figure 2: Spectrum of absorption of Curcuma extract.




Figure 3: The calibration line of Chamomile.

Figure 4: The calibration line of Turmeric.




Figure 5 : Gallic acid calibration curve.




Figure 6: Flavonoid calibration curve.




Figure 7: Tannins calibration curve.

## ANNEXE

### 1. Results of antibacterial activity of extracts:


**Table 1:** Diameter of inhibition zones of extracts by disk methods.

| The different strains | Aqueous extracts |     |     |
|-----------------------|------------------|-----|-----|
|                       | Thy              | Cam | Cur |
| <i>E-Coli</i>         | 11               | 11  | 0   |
| <i>Staph</i>          | 10               | 13  | 0   |
| <i>Pseudo</i>         | 11               | 12  | 0   |
| <i>Candida</i>        | 0                | 12  | 7   |

### 2. Results of antibacterial activity of creams:


**Table 2:** Diameter of inhibition zones of creams by well method.

| The different strains | Creams |      |    |      |    |      |    |      |    |      |    |      |    |      |
|-----------------------|--------|------|----|------|----|------|----|------|----|------|----|------|----|------|
|                       | L1     | L1.0 | L2 | L2.0 | L3 | L3.0 | L4 | L4.0 | L5 | L5.0 | L6 | L6.0 | L7 | L7.0 |
| <i>E-Coli</i>         | 14     | 12   | 13 | 0    | 10 | 10   | 0  | 14   | 16 | 15   | 16 | 17   | 12 | 0    |
| <i>Staph</i>          | 14     | 9    | 10 | 0    | 13 | 0    | 9  | 10   | 13 | 12   | 14 | 13   | 0  | 0    |
| <i>Pseudo</i>         | 0      | 0    | 12 | 0    | 19 | 15   | 16 | 14   | 12 | 0    | 13 | 10   | 17 | 0    |
| <i>Candida</i>        | 0      | 15   | 15 | 0    | 10 | 0    | 10 | 0    | 12 | 9    | 11 | 16   | 0  | 23   |



**Figure 8:** the results of the antibacterial tests for the ointments.


3. MIC result for the Curcumin:



**Figure 9:** MIC results for Curcumin extract.



البطاقة التكنولوجية للمشروع

|   |   |
|---|---|
| <p>عبد الرحيم ايمان / بن عودة<br/>شهيناز هيبه</p> <p>Abd Errahim Imene /<br/>Benaouda Chahinez Hiba</p>                     | <p>الاسم واللقب</p> <p>Votre prénom et nom</p> <p>Your first and last Name</p>                |
| <p>سي بيو</p> <p>CI Bio</p> <p>CI Bio</p>  | <p>الاسم التجاري<br/>للمشروع</p> <p>Intitulé de votre projet</p> <p>Title of your Project</p> |
| <p>Sarl la société à responsabilité limitée</p> <p>LLC (limited liability corporation)</p>                                  | <p>الصفة القانونية للمشروع</p> <p>Votre statut juridique</p> <p>Your legal status</p>         |
| <p>0675206380/0554991961</p>  | <p>رقم الهاتف</p> <p>Votre numéro de téléphone</p> <p>Your phone number</p>                   |
| <p><a href="mailto:/abderrahimiman23@gmail.com">/abderrahimiman23@gmail.com</a></p> <p>bnchahinez11@gmail.com</p>           | <p>البريد الالكتروني</p> <p>Votre adresse e-mail</p>  |





|                                      |   |
|--------------------------------------|---|
|                                      | Your email address  |
| بلدية تيارت - ولاية تيارت - الجزائر. | مقر مزاولة النشاط (الولاية - البلدية)<br>Votre ville ou commune d'activité<br>Your city or municipality of activity |

**m**

### طبيعة المشروع

|   |  |
|---|--|
| <p>في وقتنا الحالي اصبح الحصول على منتجات طبيعية فعالة في علاج شتى الافات الصحية شيئا شبه منعدم وتحت هذا الايطار تدخل منتجاتنا كحل فعال وبديل للمنتجات المعروضة في السوق من مرعم و كريم أساس و شامبو فنحن نقدم لزبائننا من نساء ورجال مرهم ضد الالتهابات للاستعمال الخارجي بسعر رمزي 550 دينار جزائري بالإضافة الى مجموعة للعناية بالشعر تتكون من شامبو و كريم أساس للاستعمال على الشعر بسعر مناسب للزبون 1500 دينار جزائري</p> <p>مقر مزاولة التصنيع ببلدية تيارت - ولاية تيارت - الجزائر.</p> | <p>مؤسسة ناشئة تهتم بصناعة المواد الصيدلانية بالإضافة الى بعض مواد التجميل.</p> <p>مؤسسة ناشئة تعتمد في صناعاتها على مواد طبيعية كالنباتات الطبية.</p> |
|---|--|

القيمة المقترحة او العرض المقترح Value Proposition

تحديد المشكل الذي يواجهه الزبون.



|   |  |
|---|--|
| <p>في الآونة الأخيرة طغت المواد الكيميائية على منتجاتنا لما أصبحت تسببه من اعراض جانبية تعود بالضرر مع مرور الوقت على المستهلك.</p>   | <p>ماهي المشكلة التي تريد حلها؟</p>                                      |
| <p>طور الدراسات</p>   | <p>ماهي البيانات المتوفرة لديك التي تبين المشكلة اتي تود دراستها ؟</p>   |
| <p>هناك بعض المشاريع التي استهدفت هذا المجال لكن الثمن المعروض في السوق الذي يتناسب مع الحد الأدنى للدخل لكن منتجاتنا المصنعة مع ضمانات وامان وفعالية وسهولة في الاستخدام تستهدف كل الفئات وهذا ما يضمن ان منتجاتنا ستكون مربحة عموم وستحل العديد من المشاكل على وجه خاص.</p> | <p>ماهي المشاريع الأخرى التي استهدفت نفس المشكلة والتي جرى تنفيذها ؟</p> |



|  |   |
|--|---|
| <p>Burt's bees</p> <p>Eminence Organique Skin Care</p> <p>Lush</p> <p>Aveda</p>  |   |
| <p>الهدف من مشروعنا الترشيد في استعمال المواد الكيميائية خاصة في مجال الصيدلة والتجميل.</p> <p>انشاء مؤسسات أخرى لكي نصل الى مجموعة شاملة متنوعة المنتجات بأسعار تنافسية في السوق وهذا إجابة الى احتياجات و متطلبات الزبائن.</p> <p>بعد الهدف المعروض سابقا نطمح الى ان تصبح منتجاتنا معروفة على المستوى الوطني والعالمي وهذا عن طريق انشاء شراكات مع الدول الأجنبية.</p> <p>وعلى المدى البعيد للمشروع سنعمل جاهدين الى توفير مجموعة خاصة بمرضى السرطان والمصابين بداء الثعلبية و ذلك بتطوير تركيبة منتجاتنا لمساندة كل فئة مرضية.</p> | <p>ماهي اهداف مشروعك و/أو نتائجه المتوقعة ؟</p> |



## القيمة المقترحة وفق المعايير التالية

|  |                            |
|--|----------------------------|
| تصنيع مواد أخرى كسيروم للشعر إضافة الى كريم للاستعمال الدائم للمحافظة على شعر صحي وناعم.   | القيمة المبتكرة او الجديدة |
| انه تقريبا كل المنتجات الطبيعية المعروضة في السوق تحتوي على نسبة معتبرة من بعض المواد الكيميائية لكن نحن نسعى الى توفير منتج طبيعي امن خالي من المضافات الكيميائية قدر المستطاع. | القيمة بالتخصيص            |
| نعرض منتجاتنا للسوق بسعر رمزي متفق عليه مقارنة بالمنتجات الأخرى<br>550 دينار جزائري لعلبة المرهم و 1500 دينار جزائري<br>لمجموعة العناية بالشعر .                                 | القيمة بالسعر              |



|  |   |
|--|---|
| <p>كريمة الالتهابات بسعة 15 غ وكريم أساس<br/>للشعر بسعة 15 غ وشامبو مرافق للشعر<br/>بحجم 100 مل يحفظ بعيدا عن اشعة الشمس<br/>مدة الصلاحية 6 أشهر.</p>            | <p>القيمة بالتصميم</p>  |
| <p>الجودة العالية والفعالية، الموثوقية والابتكار، بالإضافة الى التوافق مع<br/>متطلبات السوق.</p>   | <p>القيمة بالاداء العالي</p>  |
| <p>نقدم منتجات شاملة ذات جودة عالية تمثل أفضل<br/>اخبارات في السوق.<br/>خدمة العملاء المتميزة بالإضافة الى التوافق مع المعايير<br/>البيئية والاجتماعية.</p>      | <p>القيمة بالخدمة الشاملة</p>   |
| <p>نحن نقدم منتجات صيدلانية وتجميلية تعتمد على تقنيات جديدة<br/>ومبتكرة تهدف الى تحقيق تفوق وابتكار يمكن انا يحدث فرقا في<br/>صناعة الجمال والرعاية الشخصية.</p> | <p>القيمة المبتكرة او الجديدة</p>   |
| <p>منتوج طبيعي جزائري.<br/>مصنع نباتات معروفة لدى الزبائن.</p>   | <p>قيم أخرى</p>   |



Customer Segments شرائح العملاء او الزبائن

| Géographique                                    | Démographique (B2C) | Démographique (B2B)                          | Psychographique  | Comportementa                                  |
|---|---------------------|--|--|--|
| الجغرافية<br>الجهة الغربية ,<br>الوسطى والشرقية |                     | الموزعون<br><br>مراكز التجميل<br>مراكز البيع | العوامل النفسية و<br>الشخصية<br><br>المنتوج يعطي<br>انطباعي إيجابي لدى الزبائن | السلوكيات<br><br>سلوك<br>المجتمع<br>منتوج حلال |
| Continent                                       | Age                 | Secteur                                      | Classe sociale   | Usage  |



|   |  |   |  |   |
|---|--|---|--|---|
| القارة<br>افريقيا   | العمر 15 سنة<br>الى 50 سنة                                 | القطاع<br>الكيمياء<br>التصنيعية                           | الطبقة الاجتماعية الفئة<br>العليا<br>المتوسطة و الدنيا | استخدام لعلاج<br>تساقط الشعر<br>اثناء الاستحمام   |
| Pays<br>الدولة<br>الجزائر   | Sexe<br>الجنس<br>نساء و رجال                               | Nombre<br>d'employés<br>عدد العمال في<br>القطاع<br>6 عمال | Niveau de vie<br>المستوى<br>المعيشي                    | Loyauté<br>الوفاء<br>الاستخدام يدوم لمدة 6<br>اشهر متواصلة كل علبة<br>للمنتوج صالحة لمدة شهر  |
| Région<br>الجهة الجغرافية<br>المستهدفة<br>تيارت ثم<br>غليزان ثم العاصمة<br>للتسويق نحو كل<br>ولايات الوطن | Revenus annuel<br>متوسط الدخل<br>40000 الف دينار<br>جزائري | Maturité de<br>l'entreprise<br>نضج المؤسسة                | Valeurs<br>القيم<br>ابتكارية<br>علمية<br>ثقافية        | Intérêt<br>اهتمام<br>الجانب<br>البيسيكولوجي<br>الشعر الصحي<br>والناعم يضيفي<br>جمالية<br>واسترجاع الثقة<br>بالنفس للزبائن<br>المتضررة من مشكل |





|   |  |   |  |   |
|---|--|---|--|---|
|   |  |   |  | تساقط الشعر   |
| Département<br>الولاية<br>تيارت                     | Etat matrimonial<br>الحالة الاجتماعية                        | Situation<br>financière<br>الحالة المالية<br>للمؤسسة<br>مؤسسة ناشئة تعتمد<br>على موارد مالية<br>بسيطة | Personnalité<br>الشخصية  | Passion<br>الهوية و الشغف   |
| Ville<br>دائرة الرحوية بلدية<br>الرحوية ولاية تيارت | Niveau d'étude<br>المستوى<br>الدراسي<br>ماستر 2 كيمياء عضوية | Détention/<br>actionariat<br>الملكية/المساهمة<br>قطعة ارض خاصة  | Convictions<br>المعتقدات<br>منتوج حلال يعتمد<br>على تركيبة طبيعية<br>بنينات معروفة لدى<br>المجتمع مع بفعاليتها | Sensibilité<br>حساسيات  |
| Quartier<br>الحي<br>العنوان .....                   | Profession<br>المهنة<br>القطاع التجاري                       | Valorisation/<br>capitalisation<br>boursière<br>التقييم / القيمة                                      | Présence<br>digitale et sur<br>les réseaux<br>sociaux  | Habitude de<br>consommation<br>عادة الاستهلاك<br>مرتين في الأسبوع |



|   |   | السوقية السعر<br>الجودة                           | استعمال التكنولوجيا في<br>التواصل<br>منصة رقمية  |  |
|---|---|---|--|--|
| Climat<br>المناخ<br>يحفظ بعيدا عن<br>اشعة الشمس | Culture<br>الثقافة<br>سهولة في الترويج<br>وهذا عائد<br>الى المعرفة<br>السابقة للنباتات<br>المستعملة | Business mod<br>نموذج العمل<br>مكون من 24<br>صفحة | Centres<br>d'intérêts<br>مركز الاهتمام<br>كريمة للالتهايات<br>بسة 15 غ و<br>كزومة للشعر بسة<br>15 غ وشامبو<br>مرافق للاستعمال<br>بحجم 100 مل | Mode de<br>paiement<br>طرق الدفع<br>دفع نقدي |
|   | Religion<br>الدين<br>الإسلام<br>حلال  | Secteur servi<br>القطاع الذي يخدمه                |  | Connaissance<br>المعرفة                      |
|   | Langue<br>اللغة   | Technologie<br>utilisée                           |  | Nature de la<br>demande                      |



|  |                                   |   |   |
|--|-----------------------------------|---|---|
|  | العربية<br>الفرنسية<br>الإنجليزية | التكنولوجيا المستعملة<br>تسهيل التصنيع مع المحافظة<br>على الجودة  | طبيعة الطلب كذمة<br>للاستعمال قبل<br>الاستحمام على الشعر<br>يلبها الشامبو يستعمل من<br>طرف الأشخاص من<br>15<br>فما فوق. |
|  |                                   | ou Format du<br>produit<br>packaging<br>كريمة سهلة<br>الاستخدام بسعة<br>15 غ مع شامبو<br>مرافق بحجم 100<br>مل | Fréquence<br>d'achat<br>عدد مرات الطلب<br>على السلعة فعالية<br>المنتوج تضمن<br>استمرارية الطلب<br>عليه                  |

### قنوات التوزيع Channels

|  |                              |                   |
|--|------------------------------|-------------------|
| <p>046 25 61 33<br/>incubator@univ-tiaret.dz<br/>Ibn khaldoun University</p> | متاجر خاصة و موقع الكتروني . | المبيعات المباشرة |
|--|------------------------------|-------------------|



|               |              |
|---------------|--------------|
|               | تجار الجملة  |
| مراكز التجميل | الموزعون     |
| الموزعون      | توسع التجارة |

### العلاقة مع العملاء Customer Relationship

|   |  |
|---|--|
| <p>1. فهم مشاكل واحتياجات العملاء.</p> <p>2. توفير حلول مخصصة.</p> <p>3. الاتصال الفعال.</p> <p>4. بناء الثقة و المصداقية.</p> <p>5. استخدام التكنولوجيا.</p> <p>6. التعلم المستمر.</p> | <p>كيف تدير علاقتك مع العملاء ؟</p>  |
| <p>Salesforce</p> <p>Hup spot CRM</p> <p>Sap CRM</p>  | <p>ماهي اهم البرامج التي ستعتمد عليها في ادارة العلاقة مع الزبون</p> <p>Microsoft Dynamics</p> <p>Monday CRM</p> <p>Zoho CRM</p> <p>الخ.....</p> |



## الشركاء الاساسيون Key Partners

| طبيعة الشراكة  | معلومات حول الشركاء                               | الشركاء       |
|--|---|---------------|
| المساعدة في الجانب التسويقي للمنتجات المصنعة                 | مستشار في التسويق بوهدة محمد<br>0796214818        | الشريك الأول  |
| مع الاهتمام بجانب التعامل العملاء عن طريق المنصة الالكترونية | منصة الكترونية مستشار بوريح<br>إسماعيل 0662999611 | الشريك الثاني |
| توفير التعليب الخاص بالمنتجات                                | ناجي اومبالاج<br>0562630149                       | الشريك الثالث |

نم بكنابة قائمة الشركاء الرئيسيون لمشروعك بالتفصيل مع ذكر الاسم، الهاتف، العنوان...إلخ



هيكل التكاليف structure Costs

|  |   |
|--|---|
| 200 مليون سنتيم  | تكاليف التعريف بالمنتج أو المؤسسة<br>Frais d'établissement  |
| 30 مليون سنتيم غاز<br>10 ملايين سنتيم ماء<br>40 مليون سنتيم كهرباء | تكاليف الحصول على العدادات ( الماء- الكهرباء<br>.....)<br>Frais d'ouverture de compteurs<br>(eaux-gaz-....) |
| 100 مليون سنتيم.   | تكاليف (التدريب- برامج العالم اللذي المختصة)<br>Logiciels, formations                                       |
| 0  | Dépôt marque, brevet, modèle<br>تكاليف براءة الاختراع و الحماية الصناعية و التجارة                          |
| 0  | Droits d'entrée<br>تكاليف الحصول على التكنولوجيا او تخصيص استعمالها   |
| 0  | Achat fonds de commerce ou parts<br>شراء الأصول التجارية أو الأسهم  |
| 60 مليون سنتيم.  | Droit au bail<br>الحق في الإيجار  |
| 0  | Caution ou dépôt de garantie<br>وديعة أو وديعة تأمين  |
| 10 مليون سنتيم.  | Frais de dossier<br>رسوم إعداد الملفات  |
| 20 مليون سنتيم المحامي.<br>10 مليون سنتيم الموثق.                  | Frais de notaire ou d'avocat<br>تكاليف الموثق-المحامي-.....   |
| 100 مليون سنتيم.   | Enseigne et éléments de   |



|                  |  |
|------------------|--|
|                  | communication<br>تكاليف التعريف بالعلامة و تكاليف فترات الاتصال              |
| 0                | Achat immobilier<br>شراء العقارات  |
| 50 مليون سنتيم.  | Travaux et aménagements<br>الأعمال والتجديلات الماك                          |
| 1 مليار سنتيم.   | Matériel<br>الآلات- المركبات- الأجهزة  |
| 20 مليون سنتيم . | Matériel de bureau<br>تجهيزات المكتب   |
| 100 مليون سنتيم. | Stock de matières et produits<br>تكاليف التخون                               |
| 500 مليون سنتيم. | Trésorerie de départ<br>الندوق النودى (الندوق) الذي نحتاجه في بداية المشروع. |

المجموع = 2 مليار و 250 مليون سنتيم.

### النفقات او التكاليف الثابتة الخاصة بمشروعك

|                  |  |
|------------------|--|
| 100 مليون سنتيم. | Assurances<br>التأمينات                        |
| 15 مليون سنتيم.  | Téléphone, internet<br>الهاتف و الانترنت       |
| 30 مليون سنتيم.  | Autres abonnements<br>اشترابات أخرى            |
| 50 مليون سنتيم.  | Carburant, transports<br>الوقود و تكاليف النقل |
| 50 مليون سنتيم.  | Frais de déplacement et hébergement            |





|                  | تكاليف التنقل و المبيت   |
|------------------|--|
| 100 مليون سنتيم. | Eau, électricité, gaz<br>فواتير الماء - الكهرباء - الغاز         |
| 20 مليون سنتيم.  | Mutuelle<br>التعاضدية الاجتماعية                                 |
| 50 مليون سنتيم.  | Fournitures diverses<br>لوازم متنوعة                             |
| 50 مليون سنتيم.  | Entretien matériel et vêtements<br>صيانة المعدات و الملابس       |
| /                | Nettoyage des locaux<br>تنظيف المباني                            |
| 50 مليون سنتيم.  | Budget publicité et communication<br>ميزانية الإعلان و الاتصالات |

المجموع = 515 مليون سنتيم.

### Revenue Stream مصادر الواردات

|   |  |
|---|--|
| / | Apport personnel ou familial<br>المساهمة الشخصية أو العائلية |
| / | Apports en nature (en valeur)<br>التبرعات العينية            |
| / | Prêt n°1 (nom de la banque)<br>قرض رقم 1 اسم البنك           |
| / | Prêt n°2 (nom de la banque)<br>قرض رقم 2 اسم البنك           |
| / | Prêt n°3 (nom de la banque)<br>قرض رقم 3 اسم البنك           |



|   |  |
|---|--|
| / | Subvention n°1 (libellé)<br>منحة 1       |
| / | Subvention n°2 (libellé)<br>منحة 2       |
| / | Autre financement (libellé)<br>تمويل آخر |

المجموع = 00

رقم الأعمال

بيع المنتج في السنة الأولى **Votre chiffre d'affaires de la première année**

| متوسط أيام العمل في الشهر | بيع المنتج في السنة الأولى |
|---------------------------|----------------------------|
| 22                        | 1Mois الشهر                |
| 22                        | 2Mois الشهر                |
| 22                        | 3Mois الشهر                |
| 22                        | 4Mois الشهر                |
| 22                        | 5Mois الشهر                |
| 22                        | 6Mois الشهر                |
| 22                        | 7Mois الشهر                |
| 22                        | 8Mois الشهر                |
| 22                        | 9Mois الشهر                |
| 22                        | 10Mois الشهر               |
| 22                        | 11Mois الشهر               |
| 22                        | 12Mois الشهر               |

المجموع = 264 يوم سبوتاً.



النسبة المئوية للزيادة في حجم الأعمال بين كل شهر لسنة الأولى؟ 10%

**Votre chiffre d'affaires de la deuxième année** بيع المنتج في السنة الثانية

| متوسط أيام العمل في الشهر | بيع المنتج في السنة الثانية |
|---------------------------|-----------------------------|
| 23                        | 1Mois الشهر                 |
| 23                        | 2Mois الشهر                 |
| 23                        | 3Mois الشهر                 |
| 23                        | 4Mois الشهر                 |
| 23                        | 5Mois الشهر                 |
| 23                        | 6Mois الشهر                 |
| 23                        | 7Mois الشهر                 |
| 23                        | 8Mois الشهر                 |
| 23                        | 9Mois الشهر                 |
| 23                        | 10Mois الشهر                |
| 23                        | 11Mois الشهر                |
| 23                        | 12Mois الشهر                |

المجموع = 276 يوم سبوتيا.

النسبة المئوية للزيادة في حجم الأعمال بين كل شهر لسنة الثانية؟ 15%

**Votre chiffre affaires de la troisième année** بيع المنتج في السنة الثالثة

| متوسط أيام العمل في الشهر | بيع المنتج في السنة الثالثة |
|---------------------------|-----------------------------|
| 24                        | 1Mois الشهر                 |
| 24                        | 2Mois الشهر                 |
| 24                        | 3Mois الشهر                 |
| 24                        | 4Mois الشهر                 |



|    |              |
|----|--------------|
| 24 | 5Mois الشهر  |
| 24 | 6Mois الشهر  |
| 24 | 7Mois الشهر  |
| 24 | 8Mois الشهر  |
| 24 | 9Mois الشهر  |
| 24 | 10Mois الشهر |
| 24 | 11Mois الشهر |
| 24 | 12Mois الشهر |

المجموع = 288 يوم سنوياً.

النسبة المئوية للزيادة في حجم الأعمال بين كل شهر لسنة الثالثة؟ 25%

تطور حجم رزم الأعمال ني السنة

- النسبة المئوية للزيادة في حجم الأعمال بين السنة 1 والسنة 2؟ 5%
- النسبة المئوية للزيادة في حجم الأعمال بين السنة 2 والسنة 3؟ 10%

حاجتك لرأس المال العام

|        |  |
|--------|--|
| 30 يوم | متوسط مدة الائتمانات الممنوحة للعملاء بالأيام<br>Durée moyenne des crédits accordés aux clients en jours |
| 30 يوم | متوسط مدة ديون الموردين بالأيام<br>Durée moyenne des dettes fournisseurs en jours                        |

رواتب الموظفين و مسؤولين الشركة

|              |                                     |
|--------------|-------------------------------------|
| 30000 / شهر. | رواتب الموظفين<br>Salaires employés |
| 60000 / شهر. | صافي أجور المسؤولين                 |



Rémunération nette dirigeant

# Business Model Canvas

Designed for:

Designed by:

Date:

Version:

## Key Partners

Who are our Key Partners? Who are our key suppliers? Which Key Resources are we acquiring from partners? Which Activities do partners perform?

MOTIVATIONS FOR PARTNERS: Cost Optimization and economy, Reduction of risk and uncertainty, Acquisition of particular resources and activities

## Key Activities

What Key Activities do our Value Propositions require? Our Distribution Channels? Customer Relationships? Revenue Streams?

### CATEGORIES:

Production, Problem Solving, Platform/Network

## Key Resources

What Key Resources do our Value Propositions require? Our Distribution Channels? Customer Relationships? Revenue Streams?

TYPES OF RESOURCES: Physical, Intellectual (brand patents, copyright), Human, Financial

## Value Propositions

What value do we deliver to the customer? Which one of our customer problems are we helping to solve? Which bundles of products and services are we offering to each Customer Segment? Which customer needs are we satisfying?

CHARACTERISTICS: Newness, Performance, Customization, "the Job Done", Design, Brand/Price, Cost Reduction, Risk Reduction, Accessibility, Convenience/Usage

## Customer Relationships

What type of relationship does each Customer Segment expect us to establish and maintain with them? Which ones are we established? How are they interacting with the rest of our business model? How costly are they?

## Channels

Through which Channels do our Customer Segments want to be reached? How are we reaching them now? How are our Channels integrated? Which ones are best? Which ones are most cost-effective? How are we integrating them with our routines?

## Customer Segments

For whom are we creating value? Who are our most important customers? Is our customer base a Mass Market, Niche Market, Segmented, Diversified, Multisided Platform?

## Cost Structure

What are the most important costs inherent in our business model? Which Key Resources are most expensive? Which Key Activities are most expensive?

BUSINESS MODELS: Cost Driven (leanest cost structure, low price value proposition, mass automation, extensive outsourcing), Value Driven (focused on value creation, premium proposition).

### SAMPLES:

CHARACTERISTICS: Fixed Costs (salaries, rents, utilities), Variable costs, Economies of scale, Economies of scope

## Revenue Streams

For what value are our customers really willing to pay? For what are they currently paying? How would they prefer to pay? How much does each Revenue Stream contribute to overall revenues?

### TYPES:

Transaction, Usage fee, Subscription Fees, Lending/Renting/Leasing, Licensing, Brokerage fees, Advertising

FIXED PRICING: List Price, Product feature dependent, Customer segment dependent

DYNAMIC PRICING: Negotiation (bargaining), Yield Management, Real-time-Market

# Business Model Canvas

Designed for:

Designed by:

Date:

Version:

## Key Partners

- Suppliers of medicinal plants.
- Distribution partners (such as shipping and Delivery companies).
- Health and beauty research centers.
- Pharmacies and beauty stores as distribution partners.

## Key Activities

- Research and development to create new products and improve existing ones.
- Manufacturing and packaging products.
- Marketing and promoting products.
- Managing sales and customer service.

## Key Resources

- Research and development team.
- Raw materials (medicinal plants).
- Factory or production unit.
- E-commerce and digital marketing platforms.
- Customer service and marketing staff.

## Value Propositions

- Providing safe and effective pharmaceutical and cosmetic products based on medicinal plant
  - Offering natural alternatives to traditional chemical products.
- Promoting health and beauty in a natural and sustainable way.

## Customer Relationships

- Excellent customer service and continuous support.
- Loyalty programs and special discounts for regular customers.
- Providing health and beauty tips and advice online.

## Channels




- Retail stores (pharmacies, beauty shops).
- Online sales through a company website.
- social media for marketing promotion.
- Health and beauty fairs

## Customer Segments

- General public, especially those interested in natural and healthy products.
- Pharmacies and beauty stores.
- Health institutions (such as hospitals and clinics).

أنواع المنتجات الطبيعية  
للشخص الأصحاء





**Cost Structure**

- Costs of raw materials (medicinal plants). شحال
- Production costs (factory, machinery, labor).
- Marketing and promotion costs.
- Shipping and distribution costs.
- Research and development costs .

The total 2 billion and 250 millions

**Revenue Streams**

- Selling pharmaceutical and cosmetic products. الأشخااص
- Monthly or yearly subscriptions for new or discounted products.
- Partnerships with beauty stores and pharmacies for wholesale sales.